Panorama of Dynamics and Geometry of Moduli Spaces and Applications

Lecture 2. Magic Wand Theorem

Anton Zorich University Paris Cité

YMSC, Tsinghua University, April 7, 2022

- Diffeomorphisms of surfaces
- Closed horocycle in the moduli space of tori
- Pseudo-Anosov
- diffeomorphisms
- Closed geodesics in the space of tori
- Dynamics in the moduli spaces
- Magic Wand Theorem
- Idea of Renormalization
- Solution of the windtree _problem

Diffeomorphisms of surfaces

Observation 1. Surfaces can wrap around themselves.

Cut a torus along a horizontal circle.



Observation 1. Surfaces can wrap around themselves.

Dehn twist twists progressively horizontal circles up to a complete turn on the opposite boundary component of the cylinder and then identifies the components.



Observation 1. Surfaces can wrap around themselves.

Dehn twist twists progressively horizontal circles up to a complete turn on the opposite boundary component of the cylinder and then identifies the components.



Dehn twist corresponds to the linear map $\hat{f}_h : \mathbb{R}^2 \to \mathbb{R}^2$ with the matrix $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$.

Observation 1. Surfaces can wrap around themselves.

Dehn twist twists progressively horizontal circles up to a complete turn on the opposite boundary component of the cylinder and then identifies the components.



 $\xrightarrow{f_h}$

 \mathbb{R}^2



Observation 1. Surfaces can wrap around themselves.

Dehn twist twists progressively horizontal circles up to a complete turn on the opposite boundary component of the cylinder and then identifies the components.



Dehn twist corresponds to the linear map $\hat{f}_h : \mathbb{R}^2 \to \mathbb{R}^2$ with the matrix $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$.

Changing the slope of the parallelogram pattern progressively we get a *closed path* in the space of flat tori.

 \mathbb{R}^2

Observation 1. Surfaces can wrap around themselves.

Dehn twist twists progressively horizontal circles up to a complete turn on the opposite boundary component of the cylinder and then identifies the components.



 $\xrightarrow{f_h}$

 \mathbb{R}^2

Dehn twist corresponds to the linear map $\hat{f}_h : \mathbb{R}^2 \to \mathbb{R}^2$ with the matrix $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$.

Observation 1. Surfaces can wrap around themselves.

Dehn twist twists progressively horizontal circles up to a complete turn on the opposite boundary component of the cylinder and then identifies the components.



 $\xrightarrow{\hat{f}_h}$

 \mathbb{R}^2

Dehn twist corresponds to the linear map $\hat{f}_h : \mathbb{R}^2 \to \mathbb{R}^2$ with the matrix $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$.

Observation 1. Surfaces can wrap around themselves.

Dehn twist twists progressively horizontal circles up to a complete turn on the opposite boundary component of the cylinder and then identifies the components.



 $\xrightarrow{f_h}$

 \mathbb{R}^2

Dehn twist corresponds to the linear map $\hat{f}_h : \mathbb{R}^2 \to \mathbb{R}^2$ with the matrix $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$.

Observation 1. Surfaces can wrap around themselves.

Dehn twist twists progressively horizontal circles up to a complete turn on the opposite boundary component of the cylinder and then identifies the components.



 $\xrightarrow{f_h}$

 \mathbb{R}^2

Dehn twist corresponds to the linear map $\hat{f}_h : \mathbb{R}^2 \to \mathbb{R}^2$ with the matrix $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$.

Observation 1. Surfaces can wrap around themselves.

Dehn twist twists progressively horizontal circles up to a complete turn on the opposite boundary component of the cylinder and then identifies the components.



 $\xrightarrow{f_h}$

 \mathbb{R}^2

Dehn twist corresponds to the linear map $\hat{f}_h : \mathbb{R}^2 \to \mathbb{R}^2$ with the matrix $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$.

Observation 1. Surfaces can wrap around themselves.

Dehn twist twists progressively horizontal circles up to a complete turn on the opposite boundary component of the cylinder and then identifies the components.



 $\xrightarrow{f_h}$

 \mathbb{R}^2

Dehn twist corresponds to the linear map $\hat{f}_h : \mathbb{R}^2 \to \mathbb{R}^2$ with the matrix $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$.

Observation 1. Surfaces can wrap around themselves.

Dehn twist twists progressively horizontal circles up to a complete turn on the opposite boundary component of the cylinder and then identifies the components.



 $\xrightarrow{f_h}$

 \mathbb{R}^2

Dehn twist corresponds to the linear map $\hat{f}_h : \mathbb{R}^2 \to \mathbb{R}^2$ with the matrix $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$.

Observation 1. Surfaces can wrap around themselves.

Dehn twist twists progressively horizontal circles up to a complete turn on the opposite boundary component of the cylinder and then identifies the components.



 $\xrightarrow{f_h}$

 \mathbb{R}^2

Dehn twist corresponds to the linear map $\hat{f}_h : \mathbb{R}^2 \to \mathbb{R}^2$ with the matrix $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$.

Observation 1. Surfaces can wrap around themselves.

Dehn twist twists progressively horizontal circles up to a complete turn on the opposite boundary component of the cylinder and then identifies the components.



 $\xrightarrow{f_h}$

 \mathbb{R}^2

 \mathbb{R}^2

Dehn twist corresponds to the linear map $\hat{f}_h : \mathbb{R}^2 \to \mathbb{R}^2$ with the matrix $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$.

Observation 1. Surfaces can wrap around themselves.

Dehn twist twists progressively horizontal circles up to a complete turn on the opposite boundary component of the cylinder and then identifies the components.





 \mathbb{R}^2



Consider a composition of two Dehn twists

$$g = f_v \circ f_h =$$





Consider a composition of two Dehn twists

$$g = f_v \circ f_h =$$





It corresponds to the integer linear map $\hat{g}: \mathbb{R}^2 \to \mathbb{R}^2$ with matrix

 $A = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \cdot \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$. Cutting and pasting appropriately the image parallelogram pattern we can check by hands that we can transform the new pattern to the initial square one.

Consider a composition of two Dehn twists

$$g = f_v \circ f_h =$$





It corresponds to the integer linear map $\hat{g} : \mathbb{R}^2 \to \mathbb{R}^2$ with matrix $A = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \cdot \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$. Cutting and pasting appropriately the image parallelogram pattern we can check by hands that we can transform the new pattern to the initial square one.



Consider a composition of two Dehn twists

$$g = f_v \circ f_h =$$





It corresponds to the integer linear map $\hat{g} : \mathbb{R}^2 \to \mathbb{R}^2$ with matrix $A = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \cdot \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$. Cutting and pasting appropriately the image parallelogram pattern we can check by hands that we can transform the new pattern to the initial square one.



Closed geodesics in the space of tori

Consider eigenvectors \vec{v}_{exp} and \vec{v}_{contr} of the linear transformation $A = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}$ corresponding to the eigenvalues $\lambda > 1$ and to $1/\lambda < 1$ respectively. Consider two transversal foliations on the original torus in directions of \vec{v}_{exp} and of \vec{v}_{contr} . We have just proved that expanding our torus \mathbb{T}^2 by factor λ in direction \vec{v}_{exp} and contracting it by the factor λ in direction \vec{v}_{contr} we get the original torus.

Consider a one-parameter family of flat tori obtained from the initial square torus by a continuous deformation expanding with a factor e^t in directions \vec{v}_{exp} and contracting with a factor e^{-t} in direction \vec{v}_{contr} . By construction such one-parameter family defines a closed curve in the space of flat tori: after the time $t_0 = \log \lambda$ it closes up and follows itself.

One can check that this closed curve is, actually, a closed geodesics in the moduli spaces of tori.

Closed geodesics in the space of tori

Consider eigenvectors \vec{v}_{exp} and \vec{v}_{contr} of the linear transformation $A = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}$ corresponding to the eigenvalues $\lambda > 1$ and to $1/\lambda < 1$ respectively. Consider two transversal foliations on the original torus in directions of \vec{v}_{exp} and of \vec{v}_{contr} . We have just proved that expanding our torus \mathbb{T}^2 by factor λ in direction \vec{v}_{exp} and contracting it by the factor λ in direction \vec{v}_{contr} we get the original torus.

Consider a one-parameter family of flat tori obtained from the initial square torus by a continuous deformation expanding with a factor e^t in directions \vec{v}_{exp} and contracting with a factor e^{-t} in direction \vec{v}_{contr} . By construction such one-parameter family defines a closed curve in the space of flat tori: after the time $t_0 = \log \lambda$ it closes up and follows itself.

One can check that this closed curve is, actually, a closed geodesics in the moduli spaces of tori.

Closed geodesics in the space of tori

Consider eigenvectors \vec{v}_{exp} and \vec{v}_{contr} of the linear transformation $A = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}$ corresponding to the eigenvalues $\lambda > 1$ and to $1/\lambda < 1$ respectively. Consider two transversal foliations on the original torus in directions of \vec{v}_{exp} and of \vec{v}_{contr} . We have just proved that expanding our torus \mathbb{T}^2 by factor λ in direction \vec{v}_{exp} and contracting it by the factor λ in direction \vec{v}_{contr} we get the original torus.

Consider a one-parameter family of flat tori obtained from the initial square torus by a continuous deformation expanding with a factor e^t in directions \vec{v}_{exp} and contracting with a factor e^{-t} in direction \vec{v}_{contr} . By construction such one-parameter family defines a closed curve in the space of flat tori: after the time $t_0 = \log \lambda$ it closes up and follows itself.

One can check that this closed curve is, actually, a closed geodesics in the moduli spaces of tori.

Dynamics in the moduli

- spaces
- From flat to complex structure
- From complex to flat structure
- Volume element
- Group action
- Masur—Veech Theorem

Magic Wand Theorem

Idea of Renormalization

Solution of the windtree problem

Dynamics in the moduli spaces

Consider the natural coordinate z in the complex plane, where lives the polygon. In this coordinate the parallel translations which we use to identify the sides of the polygon are represented as z' = z + const.

Since this correspondence is holomorphic, our flat surface S with punctured conical points inherits the complex structure. This complex structure extends to the punctured points.

Consider now a holomorphic 1-form dz in the complex plane. The coordinate z is not globally defined on the surface S. However, since the changes of local coordinates are defined as z' = z + const, we see that dz = dz'. Thus, the holomorphic 1-form dz on \mathbb{C} defines a holomorphic 1-form ω on S which in local coordinates has the form $\omega = dz$.

Consider the natural coordinate z in the complex plane, where lives the polygon. In this coordinate the parallel translations which we use to identify the sides of the polygon are represented as z' = z + const.

Since this correspondence is holomorphic, our flat surface S with punctured conical points inherits the complex structure. This complex structure extends to the punctured points.

Consider now a holomorphic 1-form dz in the complex plane. The coordinate z is not globally defined on the surface S. However, since the changes of local coordinates are defined as z' = z + const, we see that dz = dz'. Thus, the holomorphic 1-form dz on \mathbb{C} defines a holomorphic 1-form ω on S which in local coordinates has the form $\omega = dz$.

Consider the natural coordinate z in the complex plane, where lives the polygon. In this coordinate the parallel translations which we use to identify the sides of the polygon are represented as z' = z + const.

Since this correspondence is holomorphic, our flat surface S with punctured conical points inherits the complex structure. This complex structure extends to the punctured points.

Consider now a holomorphic 1-form dz in the complex plane. The coordinate z is not globally defined on the surface S. However, since the changes of local coordinates are defined as z' = z + const, we see that dz = dz'. Thus, the holomorphic 1-form dz on \mathbb{C} defines a holomorphic 1-form ω on S which in local coordinates has the form $\omega = dz$.

Consider the natural coordinate z in the complex plane, where lives the polygon. In this coordinate the parallel translations which we use to identify the sides of the polygon are represented as z' = z + const.

Since this correspondence is holomorphic, our flat surface S with punctured conical points inherits the complex structure. This complex structure extends to the punctured points.

Consider now a holomorphic 1-form dz in the complex plane. The coordinate z is not globally defined on the surface S. However, since the changes of local coordinates are defined as z' = z + const, we see that dz = dz'. Thus, the holomorphic 1-form dz on \mathbb{C} defines a holomorphic 1-form ω on S which in local coordinates has the form $\omega = dz$.
- Reciprocally a pair (Riemann surface, holomorphic 1-form) uniquely defines a flat structure: $z = \int \omega$.
- In a neighborhood of zero a holomorphic 1-form can be represented as $w^d dw$, where d is the **degree** of zero. The form ω has a zero of degree d at a conical point with cone angle $2\pi(d+1)$. Moreover, $d_1 + \cdots + d_n = 2g 2$.
- The moduli space \mathcal{H}_g of pairs (complex structure, holomorphic 1-form) is a \mathbb{C}^g -vector bundle over the moduli space \mathcal{M}_g of complex structures.
- The space \mathcal{H}_g is naturally stratified by the strata $\mathcal{H}(d_1, \ldots, d_n)$ enumerated by unordered partitions $d_1 + \cdots + d_n = 2g - 2$.
- Any holomorphic 1-forms corresponding to a fixed stratum $\mathcal{H}(d_1, \ldots, d_n)$ has exactly n zeroes P_1, \ldots, P_n of degrees d_1, \ldots, d_n .
- The vectors defining the polygon from the previous picture considered as complex numbers are the relative periods $\int_{P_i}^{P_j} \omega$ of ω , so each stratum $\mathcal{H}(d_1, \ldots, d_n)$ is modelled on the relative cohomology $H^1(S, \{P_1, \ldots, P_n\}; \mathbb{C})$ serving as period coordinates.

- Reciprocally a pair (Riemann surface, holomorphic 1-form) uniquely defines a flat structure: $z = \int \omega$.
- In a neighborhood of zero a holomorphic 1-form can be represented as $w^d dw$, where d is the **degree** of zero. The form ω has a zero of degree d at a conical point with cone angle $2\pi(d+1)$. Moreover, $d_1 + \cdots + d_n = 2q 2$.
- The moduli space \mathcal{H}_g of pairs (complex structure, holomorphic 1-form) is a \mathbb{C}^g -vector bundle over the moduli space \mathcal{M}_g of complex structures.
- The space \mathcal{H}_g is naturally stratified by the strata $\mathcal{H}(d_1, \ldots, d_n)$ enumerated by unordered partitions $d_1 + \cdots + d_n = 2g - 2$.
- Any holomorphic 1-forms corresponding to a fixed stratum $\mathcal{H}(d_1, \ldots, d_n)$ has exactly n zeroes P_1, \ldots, P_n of degrees d_1, \ldots, d_n .
- The vectors defining the polygon from the previous picture considered as complex numbers are the relative periods $\int_{P_i}^{P_j} \omega$ of ω , so each stratum $\mathcal{H}(d_1, \ldots, d_n)$ is modelled on the relative cohomology $H^1(S, \{P_1, \ldots, P_n\}; \mathbb{C})$ serving as period coordinates.

- Reciprocally a pair (Riemann surface, holomorphic 1-form) uniquely defines a flat structure: $z = \int \omega$.
- In a neighborhood of zero a holomorphic 1-form can be represented as $w^d dw$, where d is the **degree** of zero. The form ω has a zero of degree d at a conical point with cone angle $2\pi(d+1)$. Moreover, $d_1 + \cdots + d_n = 2q 2$.
- The moduli space \mathcal{H}_g of pairs (complex structure, holomorphic 1-form) is a \mathbb{C}^g -vector bundle over the moduli space \mathcal{M}_g of complex structures.
- The space \mathcal{H}_g is naturally stratified by the strata $\mathcal{H}(d_1, \ldots, d_n)$ enumerated by unordered partitions $d_1 + \cdots + d_n = 2g - 2$.
- Any holomorphic 1-forms corresponding to a fixed stratum $\mathcal{H}(d_1, \ldots, d_n)$ has exactly n zeroes P_1, \ldots, P_n of degrees d_1, \ldots, d_n .
- The vectors defining the polygon from the previous picture considered as complex numbers are the relative periods $\int_{P_i}^{P_j} \omega$ of ω , so each stratum $\mathcal{H}(d_1, \ldots, d_n)$ is modelled on the relative cohomology $H^1(S, \{P_1, \ldots, P_n\}; \mathbb{C})$ serving as period coordinates.

- Reciprocally a pair (Riemann surface, holomorphic 1-form) uniquely defines a flat structure: $z = \int \omega$.
- In a neighborhood of zero a holomorphic 1-form can be represented as $w^d dw$, where d is the **degree** of zero. The form ω has a zero of degree d at a conical point with cone angle $2\pi(d+1)$. Moreover, $d_1 + \cdots + d_n = 2q 2$.
- The moduli space \mathcal{H}_g of pairs (complex structure, holomorphic 1-form) is a \mathbb{C}^g -vector bundle over the moduli space \mathcal{M}_g of complex structures.
- The space \mathcal{H}_g is naturally stratified by the strata $\mathcal{H}(d_1, \ldots, d_n)$ enumerated by unordered partitions $d_1 + \cdots + d_n = 2g - 2$.
- Any holomorphic 1-forms corresponding to a fixed stratum $\mathcal{H}(d_1, \ldots, d_n)$ has exactly n zeroes P_1, \ldots, P_n of degrees d_1, \ldots, d_n .
- The vectors defining the polygon from the previous picture considered as complex numbers are the relative periods $\int_{P_i}^{P_j} \omega$ of ω , so each stratum $\mathcal{H}(d_1, \ldots, d_n)$ is modelled on the relative cohomology $H^1(S, \{P_1, \ldots, P_n\}; \mathbb{C})$ serving as period coordinates.

- Reciprocally a pair (Riemann surface, holomorphic 1-form) uniquely defines a flat structure: $z = \int \omega$.
- In a neighborhood of zero a holomorphic 1-form can be represented as $w^d dw$, where d is the **degree** of zero. The form ω has a zero of degree d at a conical point with cone angle $2\pi(d+1)$. Moreover, $d_1 + \cdots + d_n = 2q 2$.
- The moduli space \mathcal{H}_g of pairs (complex structure, holomorphic 1-form) is a \mathbb{C}^g -vector bundle over the moduli space \mathcal{M}_g of complex structures.
- The space \mathcal{H}_g is naturally stratified by the strata $\mathcal{H}(d_1, \ldots, d_n)$ enumerated by unordered partitions $d_1 + \cdots + d_n = 2g - 2$.
- Any holomorphic 1-forms corresponding to a fixed stratum $\mathcal{H}(d_1, \ldots, d_n)$ has exactly n zeroes P_1, \ldots, P_n of degrees d_1, \ldots, d_n .
- The vectors defining the polygon from the previous picture considered as complex numbers are the relative periods $\int_{P_i}^{P_j} \omega$ of ω , so each stratum $\mathcal{H}(d_1, \ldots, d_n)$ is modelled on the relative cohomology $H^1(S, \{P_1, \ldots, P_n\}; \mathbb{C})$ serving as period coordinates.

- Reciprocally a pair (Riemann surface, holomorphic 1-form) uniquely defines a flat structure: $z = \int \omega$.
- In a neighborhood of zero a holomorphic 1-form can be represented as $w^d dw$, where d is the **degree** of zero. The form ω has a zero of degree d at a conical point with cone angle $2\pi(d+1)$. Moreover, $d_1 + \cdots + d_n = 2q 2$.
- The moduli space \mathcal{H}_g of pairs (complex structure, holomorphic 1-form) is a \mathbb{C}^g -vector bundle over the moduli space \mathcal{M}_q of complex structures.
- The space \mathcal{H}_g is naturally stratified by the strata $\mathcal{H}(d_1, \ldots, d_n)$ enumerated by unordered partitions $d_1 + \cdots + d_n = 2g - 2$.
- Any holomorphic 1-forms corresponding to a fixed stratum $\mathcal{H}(d_1, \ldots, d_n)$ has exactly n zeroes P_1, \ldots, P_n of degrees d_1, \ldots, d_n .
- The vectors defining the polygon from the previous picture considered as complex numbers are the relative periods $\int_{P_i}^{P_j} \omega$ of ω , so each stratum $\mathcal{H}(d_1, \ldots, d_n)$ is modelled on the relative cohomology $H^1(S, \{P_1, \ldots, P_n\}; \mathbb{C})$ serving as *period coordinates*.

Volume element

Note that the vector space $H^1(S, \{P_1, \ldots, P_n\}; \mathbb{C})$ contains a natural integer lattice $H^1(S, \{P_1, \ldots, P_n\}; \mathbb{Z} \oplus \sqrt{-1}\mathbb{Z})$. Consider a linear volume element $d\nu$ normalized in such a way that the volume of the fundamental domain in this lattice equals one. Consider now the real hypersurface $\mathcal{H}_1(d_1, \ldots, d_n) \subset \mathcal{H}(d_1, \ldots, d_n)$ defined by the equation area(S) = 1. The volume element $d\nu$ can be naturally restricted to the hypersurface defining the volume element $d\nu_1$ on $\mathcal{H}_1(d_1, \ldots, d_n)$.

Theorem (H. Masur; W. A. Veech) The total volume $Vol(\mathcal{H}_1(d_1, \ldots, d_n))$ of every stratum is finite.

The Masur–Veech volumes of the first several low-dimensional strata were computed by M. Kontsevich and A. Zorich about 2000. The first efficient algorithm for evaluation of the Masur–Veech volume was found by A. Eskin and A. Okounkov. In particular, they proved that the Masur–Veech volume of any stratum always has the form $(p/q)\pi^{2g}$ where p/q is a rational number. By 2003 A. Eskin computed these rational numbers up for all strata to genus 10. By now we have much better knowledge of Masur–Veech volumes; we will discuss them in more details later in these lectures.



The subgroup $SL(2,\mathbb{R})$ of area preserving linear transformations acts on the "unit hyperboloid" $\mathcal{H}_1(d_1,\ldots,d_n)$. The diagonal subgroup $\begin{pmatrix} e^t & 0\\ 0 & e^{-t} \end{pmatrix} \subset SL(2,\mathbb{R})$ induces a natural flow on the stratum, which is called the *Teichmüller geodesic flow*.

Key Theorem (H. Masur; W. A. Veech) The action of the groups $SL(2, \mathbb{R})$ and $\begin{pmatrix} e^t & 0\\ 0 & e^{-t} \end{pmatrix}$ preserves the measure $d\nu_1$. Both actions are ergodic with respect to this measure on each connected component of every stratum $\mathcal{H}_1(d_1, \ldots, d_n)$.

Masur—Veech Theorem

Theorem of Masur and Veech claims that taking an arbitrary octagon as below we can contract it horizontally and expand vertically by the same factor e^t to get arbitrary close to, say, regular octagon.



Masur—Veech Theorem

Theorem of Masur and Veech claims that taking an arbitrary octagon as below we can contract it horizontally and expand vertically by the same factor e^t to get arbitrary close to, say, regular octagon.

There is no paradox since we are allowed to cut-and-paste!



Masur—Veech Theorem

Theorem of Masur and Veech claims that taking an arbitrary octagon as below we can contract it horizontally and expand vertically by the same factor e^t to get arbitrary close to, say, regular octagon.



The first modification of the polygon changes the flat structure while the second one just changes the way in which we unwrap the flat surface

Compute asymptotic intersection number again

Diffeomorphisms of surfaces

Dynamics in the moduli spaces

Magic Wand Theorem

• Invariant measures and orbit closures

- Fields Medal
- Breakthrough Prize
- Why the Magic Wand Theorem is astonishing

• Geometric counterpart of Ratner Theorem

Idea of Renormalization

Solution of the windtree _problem

Magic Wand Theorem

Magic Wand Theorem (A. Eskin–M. Mirzakhani–A. Mohammadi, 2014). The closure of any $SL(2, \mathbb{R})$ -orbit is a suborbifold. In period coordinates any $GL(2, \mathbb{R})$ -orbit closure is represented by a complexification of an \mathbb{R} -linear subspace.

Any ergodic $SL(2, \mathbb{R})$ -invariant measure is supported on a suborbifold. In period coordinates this suborbifold is represented by an affine subspace, and the invariant measure is just a usual affine measure on this affine subspace.

Theorem (S. Filip, 2014) Any $SL(2, \mathbb{R})$ -invariant orbifold is, actually, a complex orbifold.

Further developements (A. Eskin–C. McMullen–R. Mukamel–A. Wright). New examples of nontrivial $SL(2, \mathbb{R})$ –invariant orbifolds coming from families of "optimal billiards in quadrilaterals".

Magic Wand Theorem (A. Eskin–M. Mirzakhani–A. Mohammadi, 2014). The closure of any $SL(2, \mathbb{R})$ -orbit is a suborbifold. In period coordinates any $GL(2, \mathbb{R})$ -orbit closure is represented by a complexification of an \mathbb{R} -linear subspace.

Any ergodic $SL(2, \mathbb{R})$ -invariant measure is supported on a suborbifold. In period coordinates this suborbifold is represented by an affine subspace, and the invariant measure is just a usual affine measure on this affine subspace.

Theorem (S. Filip, 2014) Any $SL(2, \mathbb{R})$ -invariant orbifold is, actually, a complex orbifold.

Further developements (A. Eskin–C. McMullen–R. Mukamel–A. Wright). New examples of nontrivial $SL(2, \mathbb{R})$ –invariant orbifolds coming from families of "optimal billiards in quadrilaterals".

Magic Wand Theorem (A. Eskin–M. Mirzakhani–A. Mohammadi, 2014). The closure of any $SL(2, \mathbb{R})$ -orbit is a suborbifold. In period coordinates any $GL(2, \mathbb{R})$ -orbit closure is represented by a complexification of an \mathbb{R} -linear subspace.

Any ergodic $SL(2, \mathbb{R})$ -invariant measure is supported on a suborbifold. In period coordinates this suborbifold is represented by an affine subspace, and the invariant measure is just a usual affine measure on this affine subspace.

Theorem (S. Filip, 2014) Any $SL(2, \mathbb{R})$ -invariant orbifold is, actually, a complex orbifold.

Further developements (A. Eskin–C. McMullen–R. Mukamel–A. Wright). New examples of nontrivial $SL(2, \mathbb{R})$ –invariant orbifolds coming from families of "optimal billiards in quadrilaterals".

Magic Wand Theorem (A. Eskin–M. Mirzakhani–A. Mohammadi, 2014). The closure of any $SL(2, \mathbb{R})$ -orbit is a suborbifold. In period coordinates any $GL(2, \mathbb{R})$ -orbit closure is represented by a complexification of an \mathbb{R} -linear subspace.

Any ergodic $SL(2, \mathbb{R})$ -invariant measure is supported on a suborbifold. In period coordinates this suborbifold is represented by an affine subspace, and the invariant measure is just a usual affine measure on this affine subspace.

Theorem (S. Filip, 2014) Any $SL(2, \mathbb{R})$ -invariant orbifold is, actually, a complex orbifold.

Further developements (A. Eskin–C. McMullen–R. Mukamel–A. Wright). New examples of nontrivial $SL(2, \mathbb{R})$ –invariant orbifolds coming from families of "optimal billiards in quadrilaterals".

Fields Medal

At the International Congress of Mathematics in 2014 Maryam Mirzakhani has received a Fields Medal for *"for her exceptional contributions to dynamics and geometry of Riemann surfaces and their moduli spaces"* becoming the first woman to receive the Fields Medal.



Breakthrough Prize

Alex Eskin got 2020 Breakthrough Prize in Mathematics *"for revolutionary discoveries in the dynamics and geometry of moduli spaces of Abelian differentials, including the proof of the "Magic Wand Theorem" with Maryam Mirzakhani.*"



Why the Magic Wand Theorem is astonishing

For most of dynamical systems (including very nice and gentle ones) certain individual trajectories are disastrously complicated. In particular, after many iterations they might fill wired fractal sets.

For example, the map $f: x \mapsto \{2x\}$ homogeneously winding the circle $S^1 = \mathbb{R}/\mathbb{Z}$ twice around itself has orbits with orbit closures of (basically) any Hausdorff dimension between 0 and 1. The same map has infinite orbits avoiding certain arcs of the circle, etc. Even such elementary maps have certain (rare) orbits with a very bizarre behavior.

Bernoulli shift. In the binary representation of a real number $x \in [0; 1[$

$$x = \frac{n_1}{2} + \dots + \frac{n_k}{2^k} + \dotsb,$$

all the binary digits n_k are zeroes or ones. The map f acts on a sequence $(n_1, n_2, \ldots, n_k, \ldots)$ by erasing the first digit. This coding shows that we have, basically, a complete freedom in constructing orbits of f with peculiar behavior.

Why the Magic Wand Theorem is astonishing

For most of dynamical systems (including very nice and gentle ones) certain individual trajectories are disastrously complicated. In particular, after many iterations they might fill wired fractal sets.

For example, the map $f: x \mapsto \{2x\}$ homogeneously winding the circle $S^1 = \mathbb{R}/\mathbb{Z}$ twice around itself has orbits with orbit closures of (basically) any Hausdorff dimension between 0 and 1. The same map has infinite orbits avoiding certain arcs of the circle, etc. Even such elementary maps have certain (rare) orbits with a very bizarre behavior.

Bernoulli shift. In the binary representation of a real number $x \in [0; 1[$

$$x = \frac{n_1}{2} + \dots + \frac{n_k}{2^k} + \dotsb,$$

all the binary digits n_k are zeroes or ones. The map f acts on a sequence $(n_1, n_2, \ldots, n_k, \ldots)$ by erasing the first digit. This coding shows that we have, basically, a complete freedom in constructing orbits of f with peculiar behavior.

Why the Magic Wand Theorem is astonishing

For most of dynamical systems (including very nice and gentle ones) certain individual trajectories are disastrously complicated. In particular, after many iterations they might fill wired fractal sets.

For example, the map $f: x \mapsto \{2x\}$ homogeneously winding the circle $S^1 = \mathbb{R}/\mathbb{Z}$ twice around itself has orbits with orbit closures of (basically) any Hausdorff dimension between 0 and 1. The same map has infinite orbits avoiding certain arcs of the circle, etc. Even such elementary maps have certain (rare) orbits with a very bizarre behavior.

Bernoulli shift. In the binary representation of a real number $x \in [0; 1]$

$$x = \frac{n_1}{2} + \dots + \frac{n_k}{2^k} + \dotsb,$$

all the binary digits n_k are zeroes or ones. The map f acts on a sequence $(n_1, n_2, \ldots, n_k, \ldots)$ by erasing the first digit. This coding shows that we have, basically, a complete freedom in constructing orbits of f with peculiar behavior.

Geometric counterpart of Ratner Theorem

Consider one of the nicest possible dynamical systems: the geodesic flow on a closed compact Riemann surface of negative curvature. Its orbits live in the three-dimensional unit tangent bundle to the hyperbolic surface.

Geometric counterpart of Ratner Theorem

Consider one of the nicest possible dynamical systems: the geodesic flow on a closed compact Riemann surface of negative curvature. Its orbits live in the three-dimensional unit tangent bundle to the hyperbolic surface.

Folklore Theorem (H. Furstenberg versus B. Weiss). For any Riemann surface C of constant negative curvature and any real number d, such that $1 \le d \le 3$, there is a trajectory of the geodesic flow on the unit tangent bundle to C such that its closure has Hausdorff dimension d.

Geometric counterpart of Ratner Theorem

Consider one of the nicest possible dynamical systems: the geodesic flow on a closed compact Riemann surface of negative curvature. Its orbits live in the three-dimensional unit tangent bundle to the hyperbolic surface.

Folklore Theorem (H. Furstenberg versus B. Weiss). For any Riemann surface C of constant negative curvature and any real number d, such that $1 \le d \le 3$, there is a trajectory of the geodesic flow on the unit tangent bundle to C such that its closure has Hausdorff dimension d.

Situation with "geodesics" of higher dimensions is completely different.

Theorem (N. Shah). In a compact manifold of constant negative curvature, the closure of a totally geodesic, complete (immersed) submanifold of dimension at least 2 is a totally geodesic immersed submanifold.

The moduli space is **not** a homogeneous space, so a priori there were no reasons to hope for a rigidity theorem like the Magic Wand Theorem of Eskin, Mirzakhani, and Mohammadi!

"But still, my homeward way has proved too long. While we were wasting time there, old Poseidon, it almost seems, stretched and extended space."

Diffeomorphisms of surfaces

Dynamics in the moduli spaces

Magic Wand Theorem

Idea of Renormalization

- Asymptotic cycle
- Spectrum of "mean monodromy"
- Hodge bundle
- Idea of

renormalization

Solution of the windtree problem

J. Brodsky И все-таки ведущая домой дорога оказалась слишком длинной, как будто Посейдон, пока мы там теряли время, растянул пространство. И. Бродский

Idea of Renormalization

Asymptotic cycle for a torus

Consider a leaf of a measured foliation on a surface. Choose a short transversal segment X. Each time when the leaf crosses X we join the crossing point with the point x_0 along X obtaining a closed loop. Consecutive return points x_1, x_2, \ldots define a sequence of cycles c_1, c_2, \ldots .



The asymptotic cycle is defined as $\lim_{n\to\infty} \frac{c_n}{n} = c \in H_1(\mathbb{T}^2; \mathbb{R}).$

Theorem (S. Kerckhoff, H. Masur, J. Smillie, 1986.) For any flat surface directional flow in almost any direction is uniquely ergodic.

This implies that for almost any direction the asymptotic cycle exists and is the same for all points of the surface.

Asymptotic cycle for a torus

Consider a leaf of a measured foliation on a surface. Choose a short transversal segment X. Each time when the leaf crosses X we join the crossing point with the point x_0 along X obtaining a closed loop. Consecutive return points x_1, x_2, \ldots define a sequence of cycles c_1, c_2, \ldots .



The asymptotic cycle is defined as $\lim_{n\to\infty} \frac{c_n}{n} = c \in H_1(\mathbb{T}^2; \mathbb{R}).$

Theorem (S. Kerckhoff, H. Masur, J. Smillie, 1986.) For any flat surface directional flow in almost any direction is uniquely ergodic.

This implies that for almost any direction the asymptotic cycle exists and is the same for all points of the surface.

Asymptotic cycle in the pseudo-Anosov case

Consider a model case of the foliation in direction of the expanding eigenvector \vec{v}_u of the Anosov map $g: \mathbb{T}^2 \to \mathbb{T}^2$ with $Dg = A = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}$. Take a closed curve γ and apply to it k iterations of g. The images $g_*^{(k)}(c)$ of the

corresponding cycle $c = [\gamma]$ get almost collinear to the expanding eigenvector \vec{v}_u of A, and the corresponding curve $g^{(k)}(\gamma)$ closely follows our foliation.

Direction of the expanding eigenvector \vec{v}_u of A = Dg



Asymptotic cycle in the pseudo-Anosov case

Consider a model case of the foliation in direction of the expanding eigenvector \vec{v}_u of the Anosov map $g: \mathbb{T}^2 \to \mathbb{T}^2$ with $Dg = A = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}$. Take a closed curve γ and apply to it k iterations of g. The images $g_*^{(k)}(c)$ of the corresponding cycle $c = [\gamma]$ get almost collinear to the expanding eigenvector \vec{v}_u of A and \vec{v}_u is the expanding eigenvector \vec{v}_u of A and \vec{v}_u is the expanding eigenvector \vec{v}_u of \vec{v}_u is the expanding eigenvector \vec{v}_u of \vec{v}_u is the expanding eigenvector \vec{v}_u is the expanding eigenvector \vec{v}_u is the expanding eigenvector \vec{v}_u of \vec{v}_u is the expanding eigenvector \vec{v}_u is the expanding eigenvector eigenvector \vec{v}_u is the expanding eigenvector eigenvector \vec{v}_u is the expanding eigenvector eig

 \vec{v}_u of A, and the corresponding curve $g^{(k)}(\gamma)$ closely follows our foliation.

Direction of the expanding eigenvector \vec{v}_u of A = Dg



Asymptotic cycle in the pseudo-Anosov case

Consider a model case of the foliation in direction of the expanding eigenvector \vec{v}_u of the Anosov map $g: \mathbb{T}^2 \to \mathbb{T}^2$ with $Dg = A = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}$. Take a closed curve γ and apply to it k iterations of g. The images $g_*^{(k)}(c)$ of the corresponding cycle $c \neq [\gamma]$ get almost collinear to the expanding eigenvector \vec{v}_u of A, and the corresponding curve $g^{(k)}(\gamma)$ closely follows our foliation.

Direction of the expanding eigenvector \vec{v}_u of A = Dg





Spectrum of "mean monodromy"

Consider a vector bundle endowed with a flat connection over a manifold X^n . Having a flow on the base we can take a fiber of the vector bundle and transport it along a trajectory of the flow. When the trajectory comes close to the starting point we identify the fibers using the connection and we get a linear transformation $\mathcal{A}(x, 1)$ of the fiber; the next time we get a matrix $\mathcal{A}(x, 2)$, etc.

The multiplicative ergodic theorem says that when the flow is ergodic a *"matrix of mean monodromy"* along the flow

$$A_{mean} := \lim_{N \to \infty} \left(\mathcal{A}^*(x, N) \cdot \mathcal{A}(x, N) \right)^{\frac{1}{2N}}$$

is well-defined and constant for almost every starting point.

Lyapunov exponents correspond to logarithms of eigenvalues of this "matrix of mean monodromy". They measure the average growth rate of the norm of vectors of the bundle when we pull them along the flow using the connection. Lyapunov exponents are dynamical analogs of characteristic numbers of the bundle. It is known that they are responsible for the diffusion rate.

Spectrum of "mean monodromy"

Consider a vector bundle endowed with a flat connection over a manifold X^n . Having a flow on the base we can take a fiber of the vector bundle and transport it along a trajectory of the flow. When the trajectory comes close to the starting point we identify the fibers using the connection and we get a linear transformation $\mathcal{A}(x, 1)$ of the fiber; the next time we get a matrix $\mathcal{A}(x, 2)$, etc.

The multiplicative ergodic theorem says that when the flow is ergodic a *"matrix of mean monodromy"* along the flow

$$A_{mean} := \lim_{N \to \infty} \left(\mathcal{A}^*(x, N) \cdot \mathcal{A}(x, N) \right)^{\frac{1}{2N}}$$

is well-defined and constant for almost every starting point.

Lyapunov exponents correspond to logarithms of eigenvalues of this "matrix of mean monodromy". They measure the average growth rate of the norm of vectors of the bundle when we pull them along the flow using the connection. Lyapunov exponents are dynamical analogs of characteristic numbers of the bundle. It is known that they are responsible for the diffusion rate.

Hodge bundle and Gauss–Manin connection

Consider a natural vector bundle over the stratum with a fiber $H^1(S; \mathbb{R})$ over a "point" (S, ω) , called the *Hodge bundle*. It carries a canonical flat connection called *Gauss—Manin connection*: we have a lattice $H^1(S; \mathbb{Z})$ in each fiber, which tells us how we can locally identify the fibers. Thus, Teichmüller flow on $\mathcal{H}_1(d_1, \ldots, d_n)$ defines Lyapunov exponents.

Hodge bundle and Gauss–Manin connection

Consider a natural vector bundle over the stratum with a fiber $H^1(S; \mathbb{R})$ over a "point" (S, ω) , called the *Hodge bundle*. It carries a canonical flat connection called *Gauss—Manin connection*: we have a lattice $H^1(S; \mathbb{Z})$ in each fiber, which tells us how we can locally identify the fibers. Thus, Teichmüller flow on $\mathcal{H}_1(d_1, \ldots, d_n)$ defines Lyapunov exponents.

Theorem (A. Eskin, M. Kontsevich, A. Z., 2014). The Lyapunov exponents λ_i of the Hodge bundle $H^1_{\mathbb{R}}$ along the Teichmüller flow restricted to an $SL(2,\mathbb{R})$ -invariant suborbifold $\mathcal{L} \subseteq \mathcal{H}_1(d_1,\ldots,d_n)$ satisfy:

$$\lambda_{1} + \lambda_{2} + \dots + \lambda_{g} = \frac{1}{12} \cdot \sum_{i=1}^{n} \frac{d_{i}(d_{i}+2)}{d_{i}+1} + \sum_{\substack{\sum \\ \text{Combinatorial types} \\ \text{of flat analogs} \\ \text{of stable curves}}} \frac{\left(\text{explicit combinatorial factor}\right) \cdot \frac{\prod_{j=1}^{k} \text{Vol} \mathcal{H}_{1}(\text{adjacent simpler strata})}{\text{Vol} \mathcal{H}_{1}(d_{1}, \dots, d_{n})}$$

Idea of renormalization

We have reformulated the model problem of windtree billiard in terms of intersection indices $c(T) \circ h$ and $c(T) \circ v$ of a cycle c(T) obtained by closing up a very long piece of vertical trajectory with two given cycles h and v on a given translation surface S.

Idea: apply the Teichmüller geodesic flow to S for an appropriate time t to get a flat surface $g_t S$ located very close to the original surface S. Close up the piece of Teichmüller geodesic to get an associated pseudo-Anosov diffeomorphism $f: S \to S$.

Note that g_t exponentially contracts the vertical direction. Choosing $t \simeq \log T$ we can transform the very long cycle c(T) to an ordinary integer cycle $f_*c(T)$ of length comparable to 1.

Conclusion: to compute $c(T) \circ h = f_*c(T) \circ f_*h$ we have to figure out how the pseudo-Anosov diffeomorphism f corresponding to a very long piece of a Teichmüller geodesic twists the distinguished cycles h and v. In other words, we have to compute the *Lyapunov exponents* for the cycles cycles h and v. Diffeomorphisms of surfaces

Dynamics in the moduli spaces

Magic Wand Theorem

Idea of Renormalization

Solution of the windtree problem

• Solution of the windtree problem

• Changing the shape of the obstacle

• Removing obstacles

• Generic windtree model of high complexity

• Hyperbolic, elliptic and parabolic dynamics

• Braque's billiard

Solution of the windtree problem
Theorem (J. Chaika–A. Eskin, 2014). For any flat surface S almost all vertical directions define a Lyapunov-generic point in the orbit closure $\overline{SL(2, \mathbb{R}) \cdot S}$.

Schematic solution of a generalized windtree problem

1. Find the family of flat surfaces \mathcal{B} associated to the original family of rational billiards;

2. Find the orbit closure $\mathcal{L} = SL(2, \mathbb{R}) \cdot \mathcal{B}$ of \mathcal{B} inside the ambient moduli space (stratum).

3. Compute or estimate the relevant Lyapunov exponents of the Hodge bundle along the Teichmüller geodesic flow on \mathcal{L} .

Currently we do not have a slightest idea on how to approach the problem when the periodic obstacles are irrational.

Theorem (J. Chaika–A. Eskin, 2014). For any flat surface S almost all vertical directions define a Lyapunov-generic point in the orbit closure $\overline{SL(2, \mathbb{R}) \cdot S}$.

Schematic solution of a generalized windtree problem

1. Find the family of flat surfaces \mathcal{B} associated to the original family of rational billiards;

2. Find the orbit closure $\mathcal{L} = SL(2, \mathbb{R}) \cdot \mathcal{B}$ of \mathcal{B} inside the ambient moduli space (stratum).

3. Compute or estimate the relevant Lyapunov exponents of the Hodge bundle along the Teichmüller geodesic flow on \mathcal{L} .

Currently we do not have a slightest idea on how to approach the problem when the periodic obstacles are irrational.

Theorem (J. Chaika–A. Eskin, 2014). For any flat surface S almost all vertical directions define a Lyapunov-generic point in the orbit closure $\overline{SL(2, \mathbb{R}) \cdot S}$.

Schematic solution of a generalized windtree problem

1. Find the family of flat surfaces \mathcal{B} associated to the original family of rational billiards;

2. Find the orbit closure $\mathcal{L} = SL(2, \mathbb{R}) \cdot \mathcal{B}$ of \mathcal{B} inside the ambient moduli space (stratum).

3. Compute or estimate the relevant Lyapunov exponents of the Hodge bundle along the Teichmüller geodesic flow on \mathcal{L} .

Currently we do not have a slightest idea on how to approach the problem when the periodic obstacles are irrational.

Theorem (J. Chaika–A. Eskin, 2014). For any flat surface S almost all vertical directions define a Lyapunov-generic point in the orbit closure $\overline{SL(2, \mathbb{R}) \cdot S}$.

Schematic solution of a generalized windtree problem

1. Find the family of flat surfaces \mathcal{B} associated to the original family of rational billiards;

2. Find the orbit closure $\mathcal{L} = SL(2, \mathbb{R}) \cdot \mathcal{B}$ of \mathcal{B} inside the ambient moduli space (stratum).

3. Compute or estimate the relevant Lyapunov exponents of the Hodge bundle along the Teichmüller geodesic flow on \mathcal{L} .

Currently we do not have a slightest idea on how to approach the problem when the periodic obstacles are irrational.

Question. What diffusion rate has a windtree billiard with "generic" (in any reasonable sense) irrational polygonal obstacles? Is it, by any chance, $\frac{1}{2}$?

Changing the shape of the obstacle

Theorem (V. Delecroix, A. Z., 2015). Changing the shape of the obstacle we get a different diffusion rate. Say, for a symmetric obstacle with 4m - 4 angles $3\pi/2$ and 4m angles $\pi/2$ the diffusion rate is



Note that once again the diffusion rate depends only on the number of the corners, but not on the (almost all) lengths of the sides, or other details of the shape of the obstacle.

Changing the shape of the obstacle

Theorem (V. Delecroix, A. Z., 2015). Changing the shape of the obstacle we get a different diffusion rate. Say, for a symmetric obstacle with 4m - 4 angles $3\pi/2$ and 4m angles $\pi/2$ the diffusion rate is

$${(2m)!!\over (2m+1)!!}~~\sim {\sqrt{\pi}\over 2\sqrt{m}}~~{
m as}~m
ightarrow\infty\,.$$



Note that once again the diffusion rate depends only on the number of the corners, but not on the (almost all) lengths of the sides, or other details of the shape of the obstacle.

How would change the diffusion rate if we remove periodically one out of four obstacles in every 2×2 group of squares?



28/31

How would change the diffusion rate if we remove periodically one out of four obstacles in every 2×2 group of squares?



And what about removing periodically two obstacles in every 2×2 group?



And what about removing periodically two obstacles in every 2×2 group?



Generic windtree model of high complexity

Theorem (Fougeron'20). The diffusion rate of a periodic billiard with $n \ge 2$ random rectangular obstacles placed as in the picture equals the top Lyapunov exponent $\lambda_1^+(Q_{n+1})$ of the Kontsevich–Zorich cocycle over the moduli space of holomorphic quadratic differentials of genus g = n + 1.



Conjectures (Zorich'98, Delecroix'15, Fougeron'19).

$$\lambda_2(\mathcal{H}(m_1,\ldots,m_n)) \to \frac{1}{2} \qquad \lambda_1^+(\mathcal{Q}(d_1,\ldots,d_n)) \to \frac{1}{2} \quad \text{as } g \to +\infty$$

uniformly for all $m_1 + \cdots + m_n = 2g - 2$ and $d_1 + \cdots + d_n = 4g - 4.$

The conjecture is confirmed by extensive computer experiments. Conceptually, it indicates that *parabolic* dynamical sys-

tems of large complexity in certain aspects mimic *hyperbolic* dynamical systems. For hyperelliptic strata $\lambda_2(\mathcal{H}_g^{hyp}) \rightarrow 1$ (Eskin–Kontsevich–Möller–Zorich + Fei Yu'18).

Time averages or ergodic dynamical systems F converge to space averages: $\frac{1}{N} (g(x) + g(F(x)) + \dots + g(F^{(N-1)}(x))) = \int_X g(x) d\mu + \text{error term},$ where the **error term** is of the order $\frac{N^{\frac{1}{2}}}{N}, \frac{\log N}{N}, \frac{N^{\alpha}}{N}$ for respectively hyperbolic, elliptic and parabolic dynamical systems: $x \stackrel{F}{\mapsto} 2x \pmod{1}$ interval exchange F

Time averages or ergodic dynamical systems F converge to space averages: $\frac{1}{N} (g(x) + g(F(x)) + \dots + g(F^{(N-1)}(x))) = \int_X g(x) d\mu + \text{error term},$ where the **error term** is of the order $\frac{N^{\frac{1}{2}}}{N}, \frac{\log N}{N}, \frac{N^{\alpha}}{N}$ for respectively hyperbolic, elliptic and parabolic dynamical systems: $x \stackrel{F}{\mapsto} 2x \pmod{1}$ interval exchange F

Time averages or ergodic dynamical systems F converge to space averages: $\frac{1}{N}(g(x) + g(F(x)) + \dots + g(F^{(N-1)}(x))) = \int_X g(x)d\mu + \text{error term},$ where the **error term** is of the order $\frac{N^{\frac{1}{2}}}{N}, \frac{\log N}{N}, \frac{N^{\alpha}}{N}$ for respectively hyperbolic, elliptic and parabolic dynamical systems: $x \stackrel{F}{\mapsto} 2x \pmod{1}$ interval exchange F



Time averages or ergodic dynamical systems F converge to space averages: $\frac{1}{N}(g(x) + g(F(x)) + \dots + g(F^{(N-1)}(x))) = \int_X g(x)d\mu + \text{error term},$ where the **error term** is of the order $\frac{N^{\frac{1}{2}}}{N}, \frac{\log N}{N}, \frac{N^{\alpha}}{N}$ for respectively hyperbolic, elliptic and parabolic dynamical systems:



Time averages or ergodic dynamical systems F converge to space averages: $\frac{1}{N}(g(x) + g(F(x)) + \dots + g(F^{(N-1)}(x))) = \int_X g(x)d\mu + \text{error term},$ where the **error term** is of the order $\frac{N^{\frac{1}{2}}}{N}, \frac{\log N}{N}, \frac{N^{\alpha}}{N}$ for respectively hyperbolic, *elliptic* and *parabolic* dynamical systems: $x \stackrel{F}{\mapsto} 2x \pmod{1}$ interval exchange F

x_0		





 $x_0 x_1$

Time averages or ergodic dynamical systems F converge to space averages: $\frac{1}{N}(g(x) + g(F(x)) + \dots + g(F^{(N-1)}(x))) = \int_X g(x)d\mu + \text{error term},$ where the **error term** is of the order $\frac{N^{\frac{1}{2}}}{N}, \frac{\log N}{N}, \frac{N^{\alpha}}{N}$ for respectively hyperbolic, elliptic and parabolic dynamical systems: $x \stackrel{F}{\mapsto} 2x \pmod{1}$ $x \stackrel{F}{\mapsto} x + \varphi \pmod{1}$ interval exchange F

 x_1

 x_0

 x_0

 x_1

Time averages or ergodic dynamical systems F converge to space averages: $\frac{1}{N}(g(x) + g(F(x)) + \dots + g(F^{(N-1)}(x))) = \int_X g(x)d\mu + \text{error term},$ where the **error term** is of the order $\frac{N^{\frac{1}{2}}}{N}, \frac{\log N}{N}, \frac{N^{\alpha}}{N}$ for respectively hyperbolic, elliptic and parabolic dynamical systems: $x \stackrel{F}{\mapsto} 2x \pmod{1}$ $x \stackrel{F}{\mapsto} x + \varphi \pmod{1}$ interval exchange F

 $x_0 x_1$

Parabolic dynamics. Lyapunov exponents. *My computer experiments of* 1993^* (motivated by the study of electron transport) indicated the **error term** with $0 < \alpha < 1$ for general interval exchanges, measured foliations, etc. I discovered that α is the second Lyapunov exponent of the Hodge bundle along the Teichmüller geodesic flow on the moduli space of Abelian differentials.

 x_1

 x_0

 x_0

This discovery was extremely influential for development of *parabolic dynamics*. Various aspects of the corresponding *"Kontsevich–Zorich conjecture"* were later proved by Forni'02, Avila–Viana'07, Eskin–Kontsevich–Zorich'14, ...

 x_1

^{*} The first person who believed in and joined these experiments was M. Kontsevich.

Time averages or ergodic dynamical systems F converge to space averages: $\frac{1}{N}(g(x) + g(F(x)) + \dots + g(F^{(N-1)}(x))) = \int_X g(x)d\mu + \text{error term},$ where the **error term** is of the order $\frac{N^{\frac{1}{2}}}{N}, \frac{\log N}{N}, \frac{N^{\alpha}}{N}$ for respectively hyperbolic, *elliptic* and *parabolic* dynamical systems: $x \stackrel{F}{\mapsto} 2x \pmod{1}$ $x \stackrel{F}{\mapsto} x + \varphi \pmod{1}$ interval exchange F





Parabolic dynamics. Lyapunov exponents. *My computer experiments of* 1993^* (motivated by the study of electron transport) indicated the **error term** with $0 < \alpha < 1$ for general interval exchanges, measured foliations, etc. I discovered that α is the second Lyapunov exponent of the Hodge bundle along the Teichmüller geodesic flow on the moduli space of Abelian differentials.

^{*} The first person who believed in and joined these experiments was M. Kontsevich.

Time averages or ergodic dynamical systems F converge to space averages: $\frac{1}{N} (g(x) + g(F(x)) + \dots + g(F^{(N-1)}(x))) = \int_X g(x) d\mu + \text{error term},$ where the **error term** is of the order $\frac{N^{\frac{1}{2}}}{N}, \frac{\log N}{N}, \frac{N^{\alpha}}{N}$ for respectively hyperbolic, elliptic and parabolic dynamical systems: $x \stackrel{F}{\mapsto} 2x \pmod{1}$ interval exchange F $I_1 \stackrel{\circ}{\to} I_2$ $I_3 \stackrel{\circ}{\to} I_4$ \downarrow_{x_0} \downarrow_{x_1} $\downarrow_$

Parabolic dynamics. Lyapunov exponents. *My computer experiments of* 1993^* (motivated by the study of electron transport) indicated the **error term** with $0 < \alpha < 1$ for general interval exchanges, measured foliations, etc. I discovered that α is the second Lyapunov exponent of the Hodge bundle along the Teichmüller geodesic flow on the moduli space of Abelian differentials.

^{*} The first person who believed in and joined these experiments was M. Kontsevich.



Parabolic dynamics. Lyapunov exponents. *My* computer experiments of 1993^* (motivated by the study of electron transport) indicated the **error term** with $0 < \alpha < 1$ for general interval exchanges, measured foliations, etc. I discovered that α is the second Lyapunov exponent of the Hodge bundle along the Teichmüller geodesic flow on the moduli space of Abelian differentials.

^{*} The first person who believed in and joined these experiments was M. Kontsevich.

Time averages or ergodic dynamical systems F converge to space averages: $\frac{1}{N}(g(x) + g(F(x)) + \dots + g(F^{(N-1)}(x))) = \int_X g(x)d\mu + \text{error term},$ where the **error term** is of the order $\frac{N^{\frac{1}{2}}}{N}, \frac{\log N}{N}, \frac{N^{\alpha}}{N}$ for respectively hyperbolic, *elliptic* and *parabolic* dynamical systems: $x \stackrel{F}{\mapsto} 2x \pmod{1}$ $x \stackrel{F}{\mapsto} x + \varphi \pmod{1}$ interval exchange F





Parabolic dynamics. Lyapunov exponents. *My computer experiments of* 1993^* (motivated by the study of electron transport) indicated the **error term** with $0 < \alpha < 1$ for general interval exchanges, measured foliations, etc. I discovered that α is the second Lyapunov exponent of the Hodge bundle along the Teichmüller geodesic flow on the moduli space of Abelian differentials.

This discovery was extremely influential for development of *parabolic dynamics*. Various aspects of the corresponding *"Kontsevich–Zorich conjecture"* were later proved by Forni'02, Avila–Viana'07, Eskin–Kontsevich–Zorich'14, ...

^{*} The first person who believed in and joined these experiments was M. Kontsevich.

Time averages or ergodic dynamical systems F converge to space averages: $\frac{1}{N}(g(x) + g(F(x)) + \dots + g(F^{(N-1)}(x))) = \int_X g(x)d\mu + \text{error term},$ where the **error term** is of the order $\frac{N^{\frac{1}{2}}}{N}, \frac{\log N}{N}, \frac{N^{\alpha}}{N}$ for respectively hyperbolic, *elliptic* and *parabolic* dynamical systems: $x \stackrel{F}{\mapsto} 2x \pmod{1}$ interval exchange F

 $x_0 x_1 x_2$

 $x_0 x_2 x_1$ $x_0 x_2$ x_1

Parabolic dynamics. Lyapunov exponents. *My computer experiments of* 1993^* (motivated by the study of electron transport) indicated the **error term** with $0 < \alpha < 1$ for general interval exchanges, measured foliations, etc. I discovered that α is the second Lyapunov exponent of the Hodge bundle along the Teichmüller geodesic flow on the moduli space of Abelian differentials.

This discovery was extremely influential for development of *parabolic dynamics*. Various aspects of the corresponding *"Kontsevich–Zorich conjecture"* were later proved by Forni'02, Avila–Viana'07, Eskin–Kontsevich–Zorich'14, ...

^{*} The first person who believed in and joined these experiments was M. Kontsevich.

Time averages or ergodic dynamical systems F converge to space averages: $\frac{1}{N}(g(x) + g(F(x)) + \dots + g(F^{(N-1)}(x))) = \int_X g(x)d\mu + \text{error term},$ where the **error term** is of the order $\frac{N^{\frac{1}{2}}}{N}, \frac{\log N}{N}, \frac{N^{\alpha}}{N}$ for respectively hyperbolic, *elliptic* and *parabolic* dynamical systems: $x \stackrel{F}{\mapsto} 2x \pmod{1}$ interval exchange F

 $x_0 x_1 x_2$

 $x_0 x_2 x_1$ $x_0 x_2$ x_1

Parabolic dynamics. Lyapunov exponents. *My computer experiments of* 1993^* (motivated by the study of electron transport) indicated the **error term** with $0 < \alpha < 1$ for general interval exchanges, measured foliations, etc. I discovered that α is the second Lyapunov exponent of the Hodge bundle along the Teichmüller geodesic flow on the moduli space of Abelian differentials.

This discovery was extremely influential for development of *parabolic dynamics*. Various aspects of the corresponding *"Kontsevich–Zorich conjecture"* were later proved by Forni'02, Avila–Viana'07, Eskin–Kontsevich–Zorich'14, ...

* The first person who believed in and joined these experiments was M. Kontsevich.

Time averages or ergodic dynamical systems F converge to space averages: $\frac{1}{N}(g(x) + g(F(x)) + \dots + g(F^{(N-1)}(x))) = \int_X g(x)d\mu + \text{error term},$ where the **error term** is of the order $\frac{N^{\frac{1}{2}}}{N}, \frac{\log N}{N}, \frac{N^{\alpha}}{N}$ for respectively hyperbolic, elliptic and parabolic dynamical systems: $x \stackrel{F}{\mapsto} 2x \pmod{1}$ interval exchange F

 $x_0x_1 x_2$



Parabolic dynamics. Lyapunov exponents. *My computer experiments of* 1993^* (motivated by the study of electron transport) indicated the **error term** with $0 < \alpha < 1$ for general interval exchanges, measured foliations, etc. I discovered that α is the second Lyapunov exponent of the Hodge bundle along the Teichmüller geodesic flow on the moduli space of Abelian differentials.

^{*} The first person who believed in and joined these experiments was M. Kontsevich.

Time averages or ergodic dynamical systems F converge to space averages: $\frac{1}{N} (g(x) + g(F(x)) + \dots + g(F^{(N-1)}(x))) = \int_X g(x) d\mu + \text{error term},$ where the **error term** is of the order $\frac{N^{\frac{1}{2}}}{N}, \frac{\log N}{N}, \frac{N^{\alpha}}{N}$ for respectively hyperbolic, elliptic and parabolic dynamical systems: $x \stackrel{F}{\mapsto} 2x \pmod{1}$ interval exchange F $I_1 \stackrel{\text{or}}{\to} I_2$ $I_2 \stackrel{\text{or}}{\to} I_3 \stackrel{\text{or}}{\to} I_4$

Parabolic dynamics. Lyapunov exponents. *My* computer experiments of 1993^* (motivated by the study of electron transport) indicated the **error term** with $0 < \alpha < 1$ for general interval exchanges, measured foliations, etc. I discovered that α is the second Lyapunov exponent of the Hodge bundle along the Teichmüller geodesic flow on the moduli space of Abelian differentials.

^{*} The first person who believed in and joined these experiments was M. Kontsevich.

Time averages or ergodic dynamical systems F converge to space averages: $\frac{1}{N}(g(x)+g(F(x))+\cdots+g(F^{(N-1)}(x))) = \int_{Y} g(x)d\mu + \operatorname{error term},$ where the *error term* is of the order $\frac{N^{\frac{1}{2}}}{N}, \frac{\log N}{N}, \frac{N^{\alpha}}{N}$ for respectively hyperbolic, elliptic and parabolic dynamical systems: $x \stackrel{F}{\mapsto} x + \varphi \pmod{1}$ $x \stackrel{F}{\mapsto} 2x \pmod{1}$ interval exchange F I_2 I_3 I_A x_{\cap} x_2 x_3 $x_0x_1x_2$ x_3 x_{\cap} x_2 x_1

Parabolic dynamics. Lyapunov exponents. *My* computer experiments of 1993^* (motivated by the study of electron transport) indicated the **error term** with $0 < \alpha < 1$ for general interval exchanges, measured foliations, etc. I discovered that α is the second Lyapunov exponent of the Hodge bundle along the Teichmüller geodesic flow on the moduli space of Abelian differentials.

^{*} The first person who believed in and joined these experiments was M. Kontsevich.

Time averages or ergodic dynamical systems F converge to space averages: $\frac{1}{N}(g(x) + g(F(x)) + \dots + g(F^{(N-1)}(x))) = \int_X g(x)d\mu + \text{error term},$ where the **error term** is of the order $\frac{N^{\frac{1}{2}}}{N}, \frac{\log N}{N}, \frac{N^{\alpha}}{N}$ for respectively hyperbolic, elliptic and parabolic dynamical systems: $x \stackrel{F}{\mapsto} 2x \pmod{1}$ $x \stackrel{F}{\mapsto} x + \varphi \pmod{1}$ interval exchange F





Parabolic dynamics. Lyapunov exponents. *My computer experiments of* 1993^* (motivated by the study of electron transport) indicated the **error term** with $0 < \alpha < 1$ for general interval exchanges, measured foliations, etc. I discovered that α is the second Lyapunov exponent of the Hodge bundle along the Teichmüller geodesic flow on the moduli space of Abelian differentials.

This discovery was extremely influential for development of *parabolic dynamics*. Various aspects of the corresponding *"Kontsevich–Zorich conjecture"* were later proved by Forni'02, Avila–Viana'07, Eskin–Kontsevich–Zorich'14, ...

* The first person who believed in and joined these experiments was M. Kontsevich.

Time averages or ergodic dynamical systems F converge to space averages: $\frac{1}{N}(g(x) + g(F(x)) + \dots + g(F^{(N-1)}(x))) = \int_X g(x)d\mu + \text{error term},$ where the **error term** is of the order $\frac{N^{\frac{1}{2}}}{N}, \frac{\log N}{N}, \frac{N^{\alpha}}{N}$ for respectively hyperbolic, *elliptic* and *parabolic* dynamical systems: $x \stackrel{F}{\mapsto} 2x \pmod{1}$ $x \stackrel{F}{\mapsto} x + \varphi \pmod{1}$ interval exchange F





Parabolic dynamics. Lyapunov exponents. *My computer experiments of* 1993^* (motivated by the study of electron transport) indicated the **error term** with $0 < \alpha < 1$ for general interval exchanges, measured foliations, etc. I discovered that α is the second Lyapunov exponent of the Hodge bundle along the Teichmüller geodesic flow on the moduli space of Abelian differentials.

This discovery was extremely influential for development of *parabolic dynamics*. Various aspects of the corresponding *"Kontsevich–Zorich conjecture"* were later proved by Forni'02, Avila–Viana'07, Eskin–Kontsevich–Zorich'14, ...

^{*} The first person who believed in and joined these experiments was M. Kontsevich.

Time averages or ergodic dynamical systems F converge to space averages: $\frac{1}{N}(g(x) + g(F(x)) + \dots + g(F^{(N-1)}(x))) = \int_X g(x)d\mu + \text{error term},$ where the **error term** is of the order $\frac{N^{\frac{1}{2}}}{N}, \frac{\log N}{N}, \frac{N^{\alpha}}{N}$ for respectively hyperbolic, elliptic and parabolic dynamical systems: $x \stackrel{F}{\mapsto} 2x \pmod{1}$ $x \stackrel{F}{\mapsto} x + \varphi \pmod{1}$ interval exchange F

 $x_0x_1x_2$ x_3

Parabolic dynamics. Lyapunov exponents. *My computer experiments of* 1993^* (motivated by the study of electron transport) indicated the **error term** with $0 < \alpha < 1$ for general interval exchanges, measured foliations, etc. I discovered that α is the second Lyapunov exponent of the Hodge bundle along the Teichmüller geodesic flow on the moduli space of Abelian differentials.

This discovery was extremely influential for development of *parabolic dynamics*. Various aspects of the corresponding *"Kontsevich–Zorich conjecture"* were later proved by Forni'02, Avila–Viana'07, Eskin–Kontsevich–Zorich'14, ...

^{*} The first person who believed in and joined these experiments was M. Kontsevich.

Time averages or ergodic dynamical systems F converge to space averages: $\frac{1}{N}(g(x) + g(F(x)) + \dots + g(F^{(N-1)}(x))) = \int_X g(x)d\mu + \text{error term},$ where the **error term** is of the order $\frac{N^{\frac{1}{2}}}{N}, \frac{\log N}{N}, \frac{N^{\alpha}}{N}$ for respectively hyperbolic, elliptic and parabolic dynamical systems: $x \stackrel{F}{\mapsto} 2x \pmod{1}$ interval exchange F





Parabolic dynamics. Lyapunov exponents. *My* computer experiments of 1993^* (motivated by the study of electron transport) indicated the **error term** with $0 < \alpha < 1$ for general interval exchanges, measured foliations, etc. I discovered that α is the second Lyapunov exponent of the Hodge bundle along the Teichmüller geodesic flow on the moduli space of Abelian differentials.

Time averages or ergodic dynamical systems F converge to space averages: $\frac{1}{N} (g(x) + g(F(x)) + \dots + g(F^{(N-1)}(x))) = \int_X g(x) d\mu + \text{error term},$ where the **error term** is of the order $\frac{N^{\frac{1}{2}}}{N}, \frac{\log N}{N}, \frac{N^{\alpha}}{N}$ for respectively hyperbolic, elliptic and parabolic dynamical systems: $x \xrightarrow{F} 2x \pmod{1}$ $x \xrightarrow{F} x + \varphi \pmod{1}$ interval exchange F $I_1 \stackrel{\circ}{\to} I_2$ $I_2 \stackrel{\circ}{\to} I_3 \stackrel{\circ}{\to} I_4$

Parabolic dynamics. Lyapunov exponents. *My computer experiments of* 1993^* (motivated by the study of electron transport) indicated the **error term** with $0 < \alpha < 1$ for general interval exchanges, measured foliations, etc. I discovered that α is the second Lyapunov exponent of the Hodge bundle along the Teichmüller geodesic flow on the moduli space of Abelian differentials.

^{*} The first person who believed in and joined these experiments was M. Kontsevich.

Time averages or ergodic dynamical systems F converge to space averages: $\frac{1}{N}(g(x)+g(F(x))+\cdots+g(F^{(N-1)}(x))) = \int_{Y} g(x)d\mu + \operatorname{error term},$ where the *error term* is of the order $\frac{N^{\frac{1}{2}}}{N}, \frac{\log N}{N}, \frac{N^{\alpha}}{N}$ for respectively hyperbolic, elliptic and parabolic dynamical systems: $x \stackrel{F}{\mapsto} x + \varphi \pmod{1}$ $x \stackrel{F}{\mapsto} 2x \pmod{1}$ interval exchange F I_2 I_A I_3 I_1 x_0 x_2 $x_4 \ x_3$ $x_0x_1 x_2 x_4 x_3$ $x_2 x_A$ x_1 x_1

Parabolic dynamics. Lyapunov exponents. *My computer experiments of* 1993^* (motivated by the study of electron transport) indicated the **error term** with $0 < \alpha < 1$ for general interval exchanges, measured foliations, etc. I discovered that α is the second Lyapunov exponent of the Hodge bundle along the Teichmüller geodesic flow on the moduli space of Abelian differentials.

^{*} The first person who believed in and joined these experiments was M. Kontsevich.

Time averages or ergodic dynamical systems F converge to space averages: $\frac{1}{N}(g(x) + g(F(x)) + \dots + g(F^{(N-1)}(x))) = \int_X g(x)d\mu + \text{error term},$ where the **error term** is of the order $\frac{N^{\frac{1}{2}}}{N}, \frac{\log N}{N}, \frac{N^{\alpha}}{N}$ for respectively hyperbolic, *elliptic* and *parabolic* dynamical systems: $x \stackrel{F}{\mapsto} 2x \pmod{1}$ interval exchange F





Parabolic dynamics. Lyapunov exponents. *My computer experiments of* 1993^* (motivated by the study of electron transport) indicated the **error term** with $0 < \alpha < 1$ for general interval exchanges, measured foliations, etc. I discovered that α is the second Lyapunov exponent of the Hodge bundle along the Teichmüller geodesic flow on the moduli space of Abelian differentials.

Time averages or ergodic dynamical systems F converge to space averages: $\frac{1}{N}(g(x) + g(F(x)) + \dots + g(F^{(N-1)}(x))) = \int_X g(x)d\mu + \text{error term},$ where the **error term** is of the order $\frac{N^{\frac{1}{2}}}{N}, \frac{\log N}{N}, \frac{N^{\alpha}}{N}$ for respectively hyperbolic, *elliptic* and *parabolic* dynamical systems: $x \stackrel{F}{\mapsto} 2x \pmod{1}$ $x \stackrel{F}{\mapsto} x + \varphi \pmod{1}$ interval exchange F





Parabolic dynamics. Lyapunov exponents. *My computer experiments of* 1993^* (motivated by the study of electron transport) indicated the **error term** with $0 < \alpha < 1$ for general interval exchanges, measured foliations, etc. I discovered that α is the second Lyapunov exponent of the Hodge bundle along the Teichmüller geodesic flow on the moduli space of Abelian differentials.

This discovery was extremely influential for development of *parabolic dynamics*. Various aspects of the corresponding *"Kontsevich–Zorich conjecture"* were later proved by Forni'02, Avila–Viana'07, Eskin–Kontsevich–Zorich'14, ...

^{*} The first person who believed in and joined these experiments was M. Kontsevich.

Time averages or ergodic dynamical systems F converge to space averages: $\frac{1}{N}(g(x) + g(F(x)) + \dots + g(F^{(N-1)}(x))) = \int_X g(x)d\mu + \text{error term},$ where the **error term** is of the order $\frac{N^{\frac{1}{2}}}{N}, \frac{\log N}{N}, \frac{N^{\alpha}}{N}$ for respectively hyperbolic, elliptic and parabolic dynamical systems: $x \stackrel{F}{\mapsto} 2x \pmod{1}$ $x \stackrel{F}{\mapsto} x + \varphi \pmod{1}$ interval exchange F





Parabolic dynamics. Lyapunov exponents. *My computer experiments of* 1993^* (motivated by the study of electron transport) indicated the **error term** with $0 < \alpha < 1$ for general interval exchanges, measured foliations, etc. I discovered that α is the second Lyapunov exponent of the Hodge bundle along the Teichmüller geodesic flow on the moduli space of Abelian differentials.

This discovery was extremely influential for development of *parabolic dynamics*. Various aspects of the corresponding *"Kontsevich–Zorich conjecture"* were later proved by Forni'02, Avila–Viana'07, Eskin–Kontsevich–Zorich'14, ...

* The first person who believed in and joined these experiments was M. Kontsevich.
Billiard in a rectangular polygon, artistic view



Georges Braque, Le Billard (1944). Centre Pompidou, Paris