

Panorama of Dynamics and Geometry of Moduli Spaces and Applications

Lecture 11. Masur-Veech volume of the moduli space of quadratic differentials, random square-tiled surfaces of large genus and random multicurves of surfaces of large genus. Part 2

Anton Zorich, University Paris Cité

(after a joint work with V. Delecroix, E. Goujard and P. Zograf
arXiv.2007.04740, to appear in Inventiones Math.)

YMSC, Tsinghua University, May 26, 2022

Notion of a random multicurve

- Reminder: space of multicurves
- Multicurves on a surface of genus two and their frequencies

Random square-tiled surfaces


Idea of the proof and further conjectures

Distribution of lengths of components of a random multicurve

Notion of a random multicurve


Reminder: space of multicurves

In train-tracks piecewise-linear coordinates, integral multicurves are represented by integer points of a conical polytope (like integral homology cycles are represented by lattice points in a vector space). Colors illustrate distinct orbits of the mapping class group. Integral multicurves are represented by lattice points on faces. Mirzakhani has proved that every orbit $\text{Mod}_{g,n} \gamma$ has strictly positive *density* which can be interpreted as a “frequency” of γ or “probability” that a multicurve taken “by random” lives in $\text{Mod}_{g,n} \gamma$.




Multicurves on a surface of genus two and their frequencies


The picture below illustrates all topological types of primitive multicurves on a surface of genus two without punctures; the fractions give frequencies of non-primitive multicurves γ having a reduced multicurve $\gamma_{reduced}$ of the corresponding type.




$$\frac{16}{63}$$




$$\frac{8}{15}$$




$$\frac{1}{9}$$



$$\frac{1}{189}$$



$$\frac{1}{45}$$



$$\frac{2}{27}$$

In genus 3 there are already 41 types of multicurves, in genus 4 there are 378 types, in genus 5 there are 4554 types and this number grows faster than exponentially when genus g grows. It becomes pointless to produce tables: we need to extract a reasonable sub-collection of most common types which ideally, carry all Thurston's measure when $g \rightarrow +\infty$.

Notion of a random multicurve

Random square-tiled surfaces

- Random integers
- Random permutations
- Shape of a random multicurve?
- Random multicurves and random square-tiled surfaces
- Shape of a random multicurve
- Weights of a random multicurve
- Number of cycles in a random permutation
- Main Theorem (informally)

Idea of the proof and further conjectures

Distribution of lengths of components of a random multicurve

Shape of a random multicurve on a surface of large genus. Shape of a random square-tiled surface of large genus.

Statistics of prime decompositions: random integer numbers

The Prime Number Theorem states that an integer number n taken randomly in a large interval $[1, N]$ is prime with asymptotic probability $\frac{\log N}{N}$.

Actually, one can tell much more about prime decomposition of a large random integer. Denote by $\omega(n)$ the number of prime divisors of an integer n counted without multiplicities. In other words, if n has prime decomposition $n = p_1^{m_1} \dots p_k^{m_k}$, let $\omega(n) = k$. By the Erdős–Kac theorem, the centered and rescaled distribution prescribed by the counting function $\omega(n)$ tends to the normal distribution:

Erdős–Kac Theorem (1939)

$$\lim_{N \rightarrow +\infty} \frac{1}{N} \text{card} \left\{ n \leq N \mid \frac{\omega(n) - \log \log N}{\sqrt{\log \log N}} \leq x \right\} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-\frac{t^2}{2}} dt.$$

The subsequent results of A. Selberg (1954) and of A. Rényi and P. Turán (1958) describe the rate of convergence.

Statistics of prime decompositions: random permutations

Denote by $K_n(\sigma)$ the number of disjoint cycles in the cycle decomposition of a permutation σ in the symmetric group S_n . Consider the uniform probability measure on S_n . A random permutation σ of n elements has exactly k cycles in its cyclic decomposition with probability $\mathbb{P}(K_n(\sigma) = k) = \frac{s(n,k)}{n!}$, where $s(n, k)$ is the unsigned Stirling number of the first kind. It is immediate to see that $\mathbb{P}(K_n(\sigma) = 1) = \frac{1}{n}$. V. L. Goncharov computed the expected value and the variance of K_n as $n \rightarrow +\infty$:


$$\mathbb{E}(K_n) = \log n + \gamma + o(1), \quad \mathbb{V}(K_n) = \log n + \gamma - \zeta(2) + o(1),$$

and proved the following central limit theorem:

Theorem (V. L. Goncharov, 1944)

$$\lim_{n \rightarrow +\infty} \frac{1}{n!} \text{card} \left\{ \sigma \in S_n \mid \frac{K_n(\sigma) - \log n}{\sqrt{\log n}} \leq x \right\} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-\frac{t^2}{2}} dt.$$

What shape has a random simple closed multicurve on a surface of large genus?





Picture from a book of Danny Calegari

Questions.

- *With what probability a random primitive multicurve γ on a surface of genus g slices the surface into $1, 2, 3, \dots$ connected components?*
- *With what probability a random multicurve $m_1\gamma_1 + m_2\gamma_2 + \dots + m_k\gamma_k$ has $k = 1, 2, \dots, 3g - 3$ primitive connected components $\gamma_1, \dots, \gamma_k$?*
- *What are the typical weights m_1, \dots, m_k ?*
- *What is the shape of a random multicurve on a surface of large genus?*

Shape of a random square-tiled surface of large genus?



Questions.

- How many singular horizontal leaves (in blue on the right picture) has a random square-tiled surface of genus g ?
- Find the probability distribution for the number $K_g(S) = 1, 2, 3, \dots, 3g - 3$ of maximal horizontal cylinders (represented by red waist curves on the left picture)
- What are the typical heights h_1, \dots, h_k of the cylinders?
- What is the shape of a random square-tiled surface of large genus?

Random multicurves and random square-tiled surfaces

Denote by $K_{g,n}(\gamma)$ the number k of components of a multicurve $\gamma = \sum_{i=1}^k m_i \gamma_i$ (counted *without* multiplicities m_i) on a surface of genus g with n cusps.

Denote by $K_{g,n}(S)$ the number of maximal horizontal cylinders in the cylinder decomposition of a square-tiled surface S of genus g with n cone-angles π

Theorem (Delecroix–Goujard–Zograf–Zorich’21.). *For any genus $g \geq 2$ and for any $k \in \mathbb{N}$, the probability $p_g(k)$ that a random multicurve γ on a surface of genus g has exactly k components counted without multiplicities coincides with the probability that a random square-tiled surface S of genus g has exactly k maximal horizontal cylinders:*

$$\mathbb{P}(K_{g,n}(\gamma) = k) = \mathbb{P}(K_{g,n}(S) = k) .$$


In other words, $K_{g,n}(\gamma)$ and $K_{g,n}(S)$, considered as random variables, determine the same probability distribution for any $g, n, 3g + n \geq 4$.

From now on we consider only hyperbolic surfaces without cusps and only square-tiled surfaces without cone-angles π (i.e. the ones corresponding to *holomorphic* quadratic differentials).

Shape of a random multicurve (random square-tiled surface) on a surface of large genus in simple words


Theorem (Delecroix–Goujard–Zograf–Zorich'20.). *With probability which tends to 1 as $g \rightarrow \infty$,*

- *The reduced multicurve $\gamma_{reduced} = \gamma_1 + \dots + \gamma_k$ associated to a random integral multicurve $\gamma = m_1\gamma_1 + \dots + m_k\gamma_k$ does not separate the surface;*
- *$\gamma_{reduced}$ has about $(\log g)/2$ components and has one of the following types:*



$0.09 \log(g)$ components

...



$0.62 \log(g)$ components

... ..

$$\mathbb{P}\left(0.09 \log g < K_g(\gamma) < 0.62 \log g\right) = 1 - O\left((\log g)^{24} g^{-1/4}\right).$$

A random square-tiled surface (without conical points of angle π) of large genus has about $\frac{\log(g)}{2}$ cylinders, and all its conical points sit at the same horizontal and at the same vertical level with probability which tends to 1 as $g \rightarrow \infty$.

Weights of a random multicurve (heights of cylinders of a random square-tiled surface)

Theorem (Delecroix–Goujard–Zograf–Zorich’19). *A random integer multicurve $m_1\gamma_1 + \dots + m_k\gamma_k$ with bounded number k of primitive components is reduced (i.e., $m_1 = \dots = m_k = 1$) with probability which tends to 1 as $g \rightarrow +\infty$. In other terms, if we consider a random square-tiled surface with at most K cylinders, the heights of all cylinders would be very likely equal to 1 for $g \gg 1$.*

Theorem (Delecroix–Goujard–Zograf–Zorich’19). *A general random integer multicurve $m_1\gamma_1 + \dots + m_k\gamma_k$ is reduced (i.e., $m_1 = \dots = m_k = 1$) with probability which tends to $\frac{\sqrt{2}}{2}$ as genus grows. More generally, all weights m_1, \dots, m_k of a random multicurve are bounded from above by an integer m with probability which tends to $\sqrt{\frac{m}{m+1}}$ as $g \rightarrow +\infty$.*

In other words, for more 70% of square-tiled surfaces of large genus, the heights of all cylinders are equal to 1.

However, the mean value of $m_1 + \dots + m_k$ is infinite in any genus g .

Number of cycles in a random permutation

Given a permutation $\sigma \in S_n$ of cycle type $(1^{\mu_1} 2^{\mu_2} \dots n^{\mu_n})$ define its *weight* as

$$w_\theta(\sigma) := \theta_1^{\mu_1} \theta_2^{\mu_2} \dots \theta_n^{\mu_n},$$

where $\theta_k = \frac{\zeta(2k)}{2}$, $k \in \mathbb{N}$. Define a probability measure on S_n by setting

$$\mathbb{P}_\theta(\sigma) := \frac{w_\theta(\sigma)}{W_\theta}, \quad \text{where} \quad W_\theta := \sum_{\sigma \in S_n} w_\theta(\sigma).$$

Measures with $\theta_k = \text{const}$, $k \in \mathbb{N}$, are called *Ewens measures*; for $\text{const} = 1$ we get the uniform measure on S_n .

Number of cycles in a random permutation

Given a permutation $\sigma \in S_n$ of cycle type $(1^{\mu_1} 2^{\mu_2} \dots n^{\mu_n})$ define its *weight* as

$$w_\theta(\sigma) := \theta_1^{\mu_1} \theta_2^{\mu_2} \dots \theta_n^{\mu_n},$$

where $\theta_k = \frac{\zeta(2k)}{2}$, $k \in \mathbb{N}$. Define a probability measure on S_n by setting

$$\mathbb{P}_\theta(\sigma) := \frac{w_\theta(\sigma)}{W_\theta}, \quad \text{where} \quad W_\theta := \sum_{\sigma \in S_n} w_\theta(\sigma).$$

Measures with $\theta_k = \text{const}$, $k \in \mathbb{N}$, are called *Ewens measures*; for $\text{const} = 1$ we get the uniform measure on S_n .

The random variable $K(\sigma)$ counting the number of disjoint cycles in the cyclic decomposition of a random permutation is very well studied (Goncharov'44, ... Hwang'94–95, ... Kowalski–Nikeghbali'10,...). The corresponding probability distribution is given by the Poisson distribution with parameter depending on n , corrected by a convolution with certain explicit function independent of n .

Using this *Mod-Poisson convergence* technique we also get a very precise description of the law for the number of cycles $K(\sigma)$ in a random permutation for our nonuniform Ewens-like measure \mathbb{P}_θ .

Probability that a random permutations has k cycles

The following Lemma identifies normalized weighted multi-variate harmonic sums as total contributions of permutations having exactly k cycles to the total sum $W_{\theta,n}$.

Lemma. *Let $\theta = \{\theta_k\}_{k \geq 1}$ be non-negative real numbers and consider the associated probability measure $\mathbb{P}_{\theta,n}$ on the symmetric group S_n for some n .*

Then

$$\frac{1}{n!} \cdot \sum_{\substack{\sigma \in S_n \\ K_n(\sigma) = k}} w_{\theta}(\sigma) = \frac{1}{k!} \cdot \sum_{i_1 + \dots + i_k = n} \frac{\theta_{i_1} \theta_{i_2} \dots \theta_{i_k}}{i_1 \dots i_k},$$

where $K_n(\sigma)$ is the number of cycles in the cycle decomposition of σ and the sum in the right hand-side is taken over positive integers i_1, \dots, i_k . In other words, we have the identity in the ring $\mathbb{Q}[[t, z]]$ of formal power series in t and z

$$\sum_{n \geq 1} \sum_{\sigma \in S_n} w_{\theta}(\sigma) t^{K_n(\sigma)} \frac{z^n}{n!} = \exp \left(t \sum_{k \geq 1} \theta_k \frac{z^k}{k} \right).$$

Main Theorem (informally)

Main Theorem (Delecroix–Goujard–Zograf–Zorich’20). As g grows, the probability distribution $\mathbb{P}(K_g = k)$ rapidly becomes, basically, indistinguishable from the distribution of the number $K_{3g-3}(\sigma)$ of disjoint cycles in a \mathbb{P}_θ -random permutation σ of $3g - 3$ elements. In particular, for any $j \in \mathbb{N}$ the difference of the j -th moments of the two distributions is of the order $O(g^{-1})$.

We have an explicit asymptotic formula for all cumulants. It gives

$$\mathbb{E}(K_g) = \frac{\log(6g - 6)}{2} + \frac{\gamma}{2} + \log 2 + o(1),$$
$$\mathbb{V}(K_g) = \frac{\log(6g - 6)}{2} + \frac{\gamma}{2} + \log 2 - \frac{3}{4}\zeta(2) + o(1),$$

where $\gamma = 0.5772\dots$ denotes the Euler–Mascheroni constant.

In practice, already for $g = 12$ the match of the graphs of the distributions is such that they are visually indistinguishable.

Mod-Poisson convergence (Hwang’94–95). For any $x > 0$ the distribution of the number of cycles of a uniformly random permutation $\sigma \in S_n$ of n elements is uniformly well-approximated in a neighborhood of $x \log n$ by the Poisson distribution with parameter $\log n + a(x)$ with an explicit correction $a(x)$.

Main Theorem (informally)

Main Theorem (Delecroix–Goujard–Zograf–Zorich’20). As g grows, the probability distribution $\mathbb{P}(K_g = k)$ rapidly becomes, basically, indistinguishable from the distribution of the number $K_{3g-3}(\sigma)$ of disjoint cycles in a \mathbb{P}_θ -random permutation σ of $3g - 3$ elements. In particular, for any $j \in \mathbb{N}$ the difference of the j -th moments of the two distributions is of the order $O(g^{-1})$.

We have an explicit asymptotic formula for all cumulants. It gives

$$\mathbb{E}(K_g) = \frac{\log(6g - 6)}{2} + \frac{\gamma}{2} + \log 2 + o(1),$$
$$\mathbb{V}(K_g) = \frac{\log(6g - 6)}{2} + \frac{\gamma}{2} + \log 2 - \frac{3}{4}\zeta(2) + o(1),$$

where $\gamma = 0.5772\dots$ denotes the Euler–Mascheroni constant.

Let $\lambda_{3g-3} = \log(6g - 6)/2$. We have uniformly in $0 \leq k \leq 1.233 \cdot \lambda_{3g-3}$

$$\mathbb{P}(K_g(\gamma) = k+1) = e^{-\lambda_{3g-3}} \cdot \frac{\lambda_{3g-3}^k}{k!} \cdot \left(\frac{\sqrt{\pi}}{2\Gamma\left(1 + \frac{k}{2\lambda_{3g-3}}\right)} + O\left(\frac{k}{(\log g)^2}\right) \right).$$

Notion of a random multicurve

Random square-tiled surfaces

Idea of the proof and further conjectures

- Idea of the proof
- Intersection numbers
- Volume polynomials
- Volume of $\mathcal{Q}_{g,n}$
- Further conjectures
- Recursive relations

Distribution of lengths of components of a random multicurve

Idea of the proof and further conjectures

Schematic idea of the proof

- Observe that square-tiled surfaces corresponding to stable graphs with more than one vertex taken together contribute only $O\left(\frac{1}{g}\right)$ to the count of all square-tiled surfaces of genus g (this conjecture of ours was proved by A. Aggarwal).
- Using large genus asymptotics for the Witten–Kontsevich correlators (conjectured by us and proved by A. Aggarwal) compute the contribution of square-tiled surfaces of genus g represented by the stable graph with exactly one vertex and with j loops. Recognize in the resulting expression the multivariate harmonic sum as in the above Lemma corresponding to parameters $\theta_k = \zeta(2k)/2$, where $k = 1, 2, \dots$.
- Apply the analytic technique developed by H. Hwang for random permutations to prove mod-Poisson convergence of the resulting distribution of the number of cycles $K_n(\sigma)$ of a random permutation σ , where “randomness” is defined using parameters $\theta_k = \zeta(2k)/2$, where $k = 1, 2, \dots$.

Intersection numbers (Witten–Kontsevich correlators)

The Deligne–Mumford compactification $\overline{\mathcal{M}}_{g,n}$ of the moduli space of smooth complex curves of genus g with n labeled marked points $P_1, \dots, P_n \in C$ is a complex orbifold of complex dimension $3g - 3 + n$.

Choose index i in $\{1, \dots, n\}$. The family of complex lines cotangent to C at the point P_i forms a holomorphic line bundle \mathcal{L}_i over $\mathcal{M}_{g,n}$ which extends to $\overline{\mathcal{M}}_{g,n}$. The first Chern class of this *tautological bundle* is denoted by $\psi_i = c_1(\mathcal{L}_i)$.

Any collection of nonnegative integers satisfying $d_1 + \dots + d_n = 3g - 3 + n$ determines a positive rational “*intersection number*” (or the “*correlator*” in the physical context):

$$\langle \tau_{d_1} \dots \tau_{d_n} \rangle_g := \int_{\overline{\mathcal{M}}_{g,n}} \psi_1^{d_1} \dots \psi_n^{d_n}.$$

The famous Witten’s conjecture claims that these numbers satisfy certain recurrence relations which are equivalent to certain differential equations on the associated generating function (“*partition function in 2-dimensional quantum gravity*”). Witten’s conjecture was proved by M. Kontsevich; alternative proofs belong to A. Okounkov and R. Pandharipande, to M. Mirzakhani, to M. Kazarian and S. Lando (and there are more).

Intersection numbers (Witten–Kontsevich correlators)

The Deligne–Mumford compactification $\overline{\mathcal{M}}_{g,n}$ of the moduli space of smooth complex curves of genus g with n labeled marked points $P_1, \dots, P_n \in C$ is a complex orbifold of complex dimension $3g - 3 + n$.

Choose index i in $\{1, \dots, n\}$. The family of complex lines cotangent to C at the point P_i forms a holomorphic line bundle \mathcal{L}_i over $\mathcal{M}_{g,n}$ which extends to $\overline{\mathcal{M}}_{g,n}$. The first Chern class of this *tautological bundle* is denoted by $\psi_i = c_1(\mathcal{L}_i)$.

Any collection of nonnegative integers satisfying $d_1 + \dots + d_n = 3g - 3 + n$ determines a positive rational “*intersection number*” (or the “*correlator*” in the physical context):

$$\langle \tau_{d_1} \dots \tau_{d_n} \rangle_g := \int_{\overline{\mathcal{M}}_{g,n}} \psi_1^{d_1} \dots \psi_n^{d_n} .$$

The famous Witten’s conjecture claims that these numbers satisfy certain recurrence relations which are equivalent to certain differential equations on the associated generating function (“*partition function in 2-dimensional quantum gravity*”). Witten’s conjecture was proved by M. Kontsevich; alternative proofs belong to A. Okounkov and R. Pandharipande, to M. Mirzakhani, to M. Kazarian and S. Lando (and there are more).

Volume polynomials

Consider the moduli space $\mathcal{M}_{g,n}$ of Riemann surfaces of genus g with n marked points. Let d_1, \dots, d_n be an ordered partition of $3g - 3 + n$ into the sum of nonnegative numbers, $d_1 + \dots + d_n = 3g - 3 + n$, let \mathbf{d} be the multiindex (d_1, \dots, d_n) and let $b^{2\mathbf{d}}$ denote $b_1^{2d_1} \dots b_n^{2d_n}$.

Define the homogeneous polynomial $N_{g,n}(b_1, \dots, b_n)$ of degree $6g - 6 + 2n$ in variables b_1, \dots, b_n :

$$N_{g,n}(b_1, \dots, b_n) := \sum_{|\mathbf{d}|=3g-3+n} c_{\mathbf{d}} b^{2\mathbf{d}},$$

where

$$c_{\mathbf{d}} := \frac{1}{2^{5g-6+2n} \mathbf{d}!} \int_{\overline{\mathcal{M}}_{g,n}} \psi_1^{d_1} \dots \psi_n^{d_n}$$

Volume polynomials

Consider the moduli space $\mathcal{M}_{g,n}$ of Riemann surfaces of genus g with n marked points. Let d_1, \dots, d_n be an ordered partition of $3g - 3 + n$ into the sum of nonnegative numbers, $d_1 + \dots + d_n = 3g - 3 + n$, let \mathbf{d} be the multiindex (d_1, \dots, d_n) and let $b^{2\mathbf{d}}$ denote $b_1^{2d_1} \dots b_n^{2d_n}$.

Define the homogeneous polynomial $N_{g,n}(b_1, \dots, b_n)$ of degree $6g - 6 + 2n$ in variables b_1, \dots, b_n :

$$N_{g,n}(b_1, \dots, b_n) := \sum_{|\mathbf{d}|=3g-3+n} c_{\mathbf{d}} b^{2\mathbf{d}},$$

where

$$c_{\mathbf{d}} := \frac{1}{2^{5g-6+2n} \mathbf{d}!} \int_{\overline{\mathcal{M}}_{g,n}} \psi_1^{d_1} \dots \psi_n^{d_n}$$

Up to a numerical factor, the polynomial $N_{g,n}(b_1, \dots, b_n)$ coincides with the top homogeneous part of the Mirzakhani's volume polynomial $V_{g,n}(b_1, \dots, b_n)$ providing the Weil–Petersson volume of the moduli space of bordered Riemann surfaces:

$$V_{g,n}^{top}(b) = 2^{2g-3+n} \cdot N_{g,n}(b).$$

Volume polynomials

Consider the moduli space $\mathcal{M}_{g,n}$ of Riemann surfaces of genus g with n marked points. Let d_1, \dots, d_n be an ordered partition of $3g - 3 + n$ into the sum of nonnegative numbers, $d_1 + \dots + d_n = 3g - 3 + n$, let \mathbf{d} be the multiindex (d_1, \dots, d_n) and let $b^{2\mathbf{d}}$ denote $b_1^{2d_1} \dots b_n^{2d_n}$.

Define the homogeneous polynomial $N_{g,n}(b_1, \dots, b_n)$ of degree $6g - 6 + 2n$ in variables b_1, \dots, b_n :

$$N_{g,n}(b_1, \dots, b_n) := \sum_{|\mathbf{d}|=3g-3+n} c_{\mathbf{d}} b^{2\mathbf{d}},$$

where

$$c_{\mathbf{d}} := \frac{1}{2^{5g-6+2n} \mathbf{d}!} \int_{\overline{\mathcal{M}}_{g,n}} \psi_1^{d_1} \dots \psi_n^{d_n}$$

Define the formal operation \mathcal{Z} on monomials as

$$\mathcal{Z} : \prod_{i=1}^n b_i^{m_i} \longmapsto \prod_{i=1}^n (m_i! \cdot \zeta(m_i + 1)),$$

and extend it to symmetric polynomials in b_i by linearity.

Volume of $\mathcal{Q}_{g,n}$

Theorem (Delecroix–Goujard–Zograf–Zorich’21). *The Masur–Veech volume $\text{Vol } \mathcal{Q}_{g,n}$ of the moduli space of meromorphic quadratic differentials with n simple poles has the following value:*

$$\text{Vol } \mathcal{Q}_{g,n} = \frac{2^{6g-5+2n} \cdot (4g - 4 + n)!}{(6g - 7 + 2n)!} \cdot \sum_{\substack{\text{Weighted graphs } \Gamma \\ \text{with } n \text{ legs}}} \frac{1}{2^{\text{Number of vertices of } \Gamma - 1}} \cdot \frac{1}{|\text{Aut } \Gamma|} \cdot \mathcal{Z} \left(\prod_{\text{Edges } e \text{ of } \Gamma} b_e \cdot \prod_{\text{Vertices of } \Gamma} N_{g_v, n_v + p_v}(\mathbf{b}_v^2, \underbrace{0, \dots, 0}_{p_v}) \right),$$

The partial sum for fixed number k of edges gives the contribution of k -cylinder square-tiled surfaces.

Volume of $\mathcal{Q}_{g,n}$

Theorem (Delecroix–Goujard–Zograf–Zorich’21). *The Masur–Veech volume $\text{Vol } \mathcal{Q}_{g,n}$ of the moduli space of meromorphic quadratic differentials with n simple poles has the following value:*

$$\text{Vol } \mathcal{Q}_{g,n} = \frac{2^{6g-5+2n} \cdot (4g - 4 + n)!}{(6g - 7 + 2n)!} \cdot \sum_{\substack{\text{Weighted graphs } \Gamma \\ \text{with } n \text{ legs}}} \frac{1}{2^{\text{Number of vertices of } \Gamma - 1}} \cdot \frac{1}{|\text{Aut } \Gamma|} \cdot \mathcal{Z} \left(\prod_{\text{Edges } e \text{ of } \Gamma} b_e \cdot \prod_{\text{Vertices of } \Gamma} N_{g_v, n_v + p_v}(\mathbf{b}_v^2, \underbrace{0, \dots, 0}_{p_v}) \right),$$

Remark. The Weil–Petersson volume of $\mathcal{M}_{g,n}$ corresponds to the *constant term* of the volume polynomial $N_{g,n}(L)$ when the lengths of all boundary components are contracted to zero. To compute the Masur–Veech volume we use the *top homogeneous parts* of volume polynomials; i.e. we use them in the opposite regime when the lengths of all boundary components tend to infinity.

Keystone underlying results and further conjectures

Our results use the Delecroix–Goujard–Zograf–Zorich’19 conjecture proved in

Theorem (Aggarwal’21). *The Masur–Veech volume of the moduli space of holomorphic quadratic differentials has the following large genus asymptotics:*

$$\text{Vol } \mathcal{Q}_g \sim \frac{4}{\pi} \cdot \left(\frac{8}{3}\right)^{4g-4} \quad \text{as } g \rightarrow +\infty.$$

Keystone underlying results and further conjectures

Our results use the Delecroix–Goujard–Zograf–Zorich’19 conjecture proved in

Theorem (Aggarwal’21). *The Masur–Veech volume of the moduli space of holomorphic quadratic differentials has the following large genus asymptotics:*

$$\text{Vol } \mathcal{Q}_g \sim \frac{4}{\pi} \cdot \left(\frac{8}{3}\right)^{4g-4} \quad \text{as } g \rightarrow +\infty.$$

The similar conjecture of Eskin–Zorich’03 on the large genus asymptotics of Masur–Veech volumes of individual strata of *Abelian* differentials is recently proved by Aggarwal’19 and by Chen–Möller–Sauvaget–Zagier’20. The analogous conjecture for *quadratic* differentials still resists:

Conjecture (ADGZZ’20). *The Masur–Veech volume of any stratum of meromorphic quadratic differentials with at most simple poles has the following large genus asymptotics (with the error term uniformly small for all partitions \mathbf{d}):*

$$\text{Vol } \mathcal{Q}(d_1, \dots, d_n) \stackrel{?}{\sim} \frac{4}{\pi} \cdot \prod_{i=1}^n \frac{2^{d_i+2}}{d_i + 2} \quad \text{as } g \rightarrow +\infty,$$

under assumption that the number of simple poles is bounded or grows much slower than the genus.

Recursive relations

Initial data: $\langle \tau_0^3 \rangle = 1, \quad \langle \tau_1 \rangle = \frac{1}{24}.$

String equation:

$$\langle \tau_0 \tau_{d_1} \cdots \tau_{d_n} \rangle_{g,n+1} = \langle \tau_{d_1-1} \cdots \tau_{d_n} \rangle_{g,n} + \cdots + \langle \tau_{d_1} \cdots \tau_{d_n-1} \rangle_{g,n}.$$

Dilaton equation:

$$\langle \tau_1 \tau_{d_1} \cdots \tau_{d_n} \rangle_{g,n+1} = (2g - 2 + n) \langle \tau_{d_1} \cdots \tau_{d_n} \rangle_{g,n}.$$

Virasoro constraints (in Dijkgraaf–Verlinde–Verlinde form; $k \geq 1$):

$$\begin{aligned} \langle \tau_{k+1} \tau_{d_1} \cdots \tau_{d_n} \rangle_g &= \frac{1}{(2k+3)!!} \left[\sum_{j=1}^n \frac{(2k+2d_j+1)!!}{(2d_j-1)!!} \langle \tau_{d_1} \cdots \tau_{d_j+k} \cdots \tau_{d_n} \rangle_g \right. \\ &\quad + \frac{1}{2} \sum_{\substack{r+s=k-1 \\ r,s \geq 0}} (2r+1)!! (2s+1)!! \langle \tau_r \tau_s \tau_{d_1} \cdots \tau_{d_n} \rangle_{g-1} \\ &\quad \left. + \frac{1}{2} \sum_{\substack{r+s=k-1 \\ r,s \geq 0}} (2r+1)!! (2s+1)!! \sum_{\{1,\dots,n\}=I \amalg J} \langle \tau_r \prod_{i \in I} \tau_{d_i} \rangle_{g'} \langle \tau_s \prod_{i \in J} \tau_{d_i} \rangle_{g-g'} \right]. \end{aligned}$$

Another Keystone result and one more conjecture

We also strongly use the uniform large genus asymptotics of ψ -classes which we conjectured in 2019 and which was proved by Aggarwal:

Theorem (Aggarwal'21). *The following **uniform** asymptotic formula is valid:*

$$\begin{aligned} \int_{\overline{\mathcal{M}}_{g,n}} \psi_1^{d_1} \cdots \psi_n^{d_n} &= \\ &= \frac{1}{24^g} \cdot \frac{(6g - 5 + 2n)!}{g! (3g - 3 + n)!} \cdot \frac{d_1! \cdots d_n!}{(2d_1 + 1)! \cdots (2d_n + 1)!} \cdot (1 + \varepsilon(\mathbf{d})), \end{aligned}$$

where $\varepsilon(\mathbf{d}) = O\left(1 + \frac{(n + \log g)^2}{g}\right)$ **uniformly** for all $n = o(\sqrt{g})$ and all partitions \mathbf{d} , $d_1 + \cdots + d_n = 3g - 3 + n$, as $g \rightarrow +\infty$.

Another Keystone result and one more conjecture

We also strongly use the uniform large genus asymptotics of ψ -classes which we conjectured in 2019 and which was proved by Aggarwal:

Theorem (Aggarwal'21). *The following **uniform** asymptotic formula is valid:*

$$\int_{\overline{\mathcal{M}}_{g,n}} \psi_1^{d_1} \cdots \psi_n^{d_n} = \frac{1}{24^g} \cdot \frac{(6g - 5 + 2n)!}{g! (3g - 3 + n)!} \cdot \frac{d_1! \cdots d_n!}{(2d_1 + 1)! \cdots (2d_n + 1)!} \cdot (1 + \varepsilon(\mathbf{d})),$$

where $\varepsilon(\mathbf{d}) = O\left(1 + \frac{(n + \log g)^2}{g}\right)$ **uniformly** for all $n = o(\sqrt{g})$ and all partitions \mathbf{d} , $d_1 + \cdots + d_n = 3g - 3 + n$, as $g \rightarrow +\infty$.

Conjecture* (Delecroix–Goujard–Zograf–Zorich). *The distribution of the number of maximal horizontal cylinders in a random Abelian square-tiled surfaces of genus g gets very well approximated by the distribution of the number of disjoint cycles in a uniformly random permutation of $4g - 3$ elements as $g \rightarrow \infty$.*

We already proved that a random square-tiled surface in a stratum \mathcal{H} has a single cylinder with probability close to $\frac{1}{\dim \mathcal{H}}$.

* About 2 years of CPU-time of two independent computer experiments with strata of genera from 40 to 10 000.

Notion of a random multicurve

Random square-tiled surfaces

Idea of the proof and further conjectures

Distribution of lengths of components of a random multicurve

- Poisson–Dirichlet process
- Distribution of lengths
- Statement for random square-tiled surfaces
- Rue des Petits-Carreux

Distribution of lengths of components of a random multicurve

Poisson–Dirichlet process

Stick breaking process. Let U_1, U_2, \dots , be i.i.d. random variables supported on $[0, 1]$ with density $\theta(1 - x)^{\theta-1}$. Take a stick of length one and chop a piece of length U_1 out of it. Chop a piece of length proportional to U_2 out of the remaining part, etc. We get a random vector

$$V = (U_1, (1 - U_1)U_2, (1 - U_1)(1 - U_2)U_3, \dots).$$

The law of V is the *Griffiths–Engen–McCloskey distribution with parameter θ* . The *Poisson–Dirichlet distribution with parameter θ* is the distribution of V^\downarrow , obtained from V by rearranging its components in the decreasing order. Both distributions are very well studied. In particular,

$$\mathbb{E}(V_j^\downarrow) = \int_0^{+\infty} \frac{(\theta E_1(x))^{j-1}}{(j-1)!} e^{-x-\theta E_1(x)} dx,$$

where $E_1(x) = \int_x^{+\infty} \frac{e^{-y}}{y} dy$.

Distribution of lengths of components of a random multicurve on a surface of large genus

Consider a random multicurve $\gamma = m_1\gamma_1 + \cdots + m_k\gamma_k$ on a hyperbolic surface $X \in \mathcal{M}_g$ and rearrange the components of the vector of weighted lengths $(m_1\ell_X(\gamma_1), \dots, m_k\ell_X(\gamma_k))$ in a decreasing order to produce a vector $\ell_X^\downarrow(\gamma)$. Normalize $\ell_X^\downarrow(\gamma)$ by $\ell_X(\gamma) = m_1\ell_X(\gamma_1) + \cdots + m_k\ell_X(\gamma_k)$.

Theorem (V. Delecroix, M. Liu, 2022). *For any X in \mathcal{M}_g and for any $j \in \mathbb{N}$ the average of $\frac{(\ell_X^\downarrow)_j}{\ell_X}$ over multicurves of bounded length gives in the limit a well-defined random variable $L_j^{(g)\downarrow}$ which depends only on j and g .*

When $g \rightarrow +\infty$, the probability distribution of $L_j^{(g)\downarrow}$ weakly converges to a limiting probability distribution V_j . The distribution V_j coincides with the limiting distribution of the normalized length of the j -th longest cycle of a non-uniformly random permutation with respect to the Evens measure with parameter $\theta = \frac{1}{2}$ on S_n as $n \rightarrow +\infty$. It is the distribution of the Poisson–Dirichlet process with parameter $\theta = \frac{1}{2}$. In particular,

$$\mathbb{E}(V_1) \approx 0.758, \quad \mathbb{E}(V_2) \approx 0.171, \quad \mathbb{E}(V_3) \approx 0.049.$$

Equivalent statement for random square-tiled surfaces

Theorem (V. Delecroix, M. Liu, 2022). *Consider the decomposition of a random square-tiled surface of genus g into maximal horizontal cylinders. Consider the vector of normalized areas of these cylinders and rearrange its components in the decreasing order. The probability distribution of the resulting random vector weakly converges to the distribution of the Poisson–Dirichlet process with parameter $\theta = \frac{1}{2}$ as g tends to ∞ .*

Restricting consideration to those random square-tiled surfaces, for which each cylinder contains at most m horizontal bands of squares, where $m = 1, 2, \dots$, one gets in the limit the very same distribution of the Poisson–Dirichlet process with parameter $\theta = \frac{1}{2}$ as g tends to ∞ .

2^e ARR^t

RUE DES
PETITS CARRÉAUX