

Combinatorics

Instructor: Prof. Jie Ma

Combinatorics 2022, Tsinghua

- This class notes will be updating throughout this course.
- The course website can be found at <https://ymsc.tsinghua.edu.cn/info/1050/2595.htm>

Contents

1	Enumeration	2
1.1	Binomial Coefficients	2
1.2	Counting Mappings	4
1.3	The Binomial Theorem	5
1.4	Inclusion and Exclusion Principle (IEP)	7
1.5	Möbius Inversion Formula	9
1.6	Generating Functions	10
1.7	Random Walks	13
1.8	Exponential Generating Functions	15
2	Basics of Graphs	19
3	Double-counting Method	21
3.1	Basics	21
3.2	Sperner's Theorem	21
3.3	Littlewood-Offord Problem	24
3.4	Turán Type Problems	25
3.5	Sperner's Lemma	27

1 Enumeration

First we give some standard notation that will be used throughout this course.

- Let n be a positive integer. We will use $[n]$ to denote the set $\{1, 2, \dots, n\}$.
- Given a set X , let $|X|$ denote the size of X , that is the number of elements contained in X .
- We use “#” to express the word “number”.
- The *factorial* of n is the product

$$n! = n \cdot (n - 1) \cdots 2 \cdot 1,$$

which can be extended to all non-negative integers by letting $0! = 1$.

1.1 Binomial Coefficients

Let X be a set of size n . Define $2^X = \{A : A \subseteq X\}$ to be the family of all subsets of X . Since the size of 2^X is equal to the number of binary vectors of length $|X|$ or the number of functions from X to $\{0, 1\}$, we have $|2^X| = 2^{|X|} = 2^n$.

Let $\binom{X}{k} = \{A : A \subseteq X, |A| = k\}$, we will use $\binom{n}{k}$ to denote $|\binom{X}{k}|$. For $n < k$, we know that $\binom{n}{k} = 0$ by definition.

Fact 1.1. For integers $n > 0$ and $0 \leq k \leq n$, we have $|\binom{X}{k}| = \binom{n}{k} = \frac{n!}{k!(n-k)!}$.

Proof. If $k = 0$, then it is clear that $|\binom{X}{0}| = |\{\emptyset\}| = 1 = \binom{n}{0}$. Now we consider $k > 0$. Let

$$(n)_k := n(n-1) \cdots (n-k+1) = \frac{n!}{(n-k)!}.$$

First we will show that number of ordered k -tuples (x_1, x_2, \dots, x_k) with distinct $x_i \in X$ is $(n)_k$. There are n choices for the first element x_1 . When x_1, \dots, x_i is chosen, there are exactly $n - i$ choices for the element x_{i+1} . So the number of ordered k -tuples (x_1, x_2, \dots, x_k) with distinct $x_i \in X$ is $(n)_k$. Since any subset $A \in \binom{X}{k}$ corresponds to $k!$ ordered k -tuples, it follows that $|\binom{X}{k}| = \frac{(n)_k}{k!} = \frac{n!}{k!(n-k)!}$. This finishes the proof. ■

Next we discuss more properties of binomial coefficients.

- Fact 1.2.** (1). $\binom{n}{k} = \binom{n}{n-k}$ for $0 \leq k \leq n$.
(2). $2^n = \sum_{0 \leq k \leq n} \binom{n}{k}$.
(3). $\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$. (*Pascal's identity*)

Proof. (1) is trivial. Since $2^{[n]} = \cup_{0 \leq k \leq n} \binom{[n]}{k}$, we see $2^n = \sum_{0 \leq k \leq n} \binom{n}{k}$, proving (2). Finally, we consider (3). Note that the first term on the right hand side $\binom{n-1}{k-1}$ is the number of k -sets containing a fixed element, while the second term $\binom{n-1}{k}$ is the number of k -sets avoiding this element. So their summation gives the total number of k -sets in $[n]$, which is $\binom{n}{k}$. This finishes the proof. ■

Pascal's triangle is a triangular array constructed by summing adjacent elements in preceding rows. By Fact 1.2 (3), in the following graph we have that the k -th element in the $n + 1$ row is $\binom{n}{k-1}$.

				1										
				1	1									
				1	2	1								
				1	3	3	1							
				1	4	6	4	1						
				1	5	10	10	5	1					
				1	6	15	20	15	6	1				
				1	7	21	35	35	21	7	1			
				1	8	28	56	70	56	28	8	1		
				1	9	36	84	126	126	84	36	9	1	
				1	10	45	120	210	252	210	120	45	10	1

Fact 1.3. The number of integer solutions (x_1, \dots, x_n) to the equation $x_1 + \dots + x_n = k$ with each $x_i \in \{0, 1\}$ is $\binom{n}{k}$.

Fact 1.4. The number of integer solutions (x_1, \dots, x_n) to the equation $x_1 + \dots + x_n = k$ with each $x_i > 0$ is $\binom{k-1}{n-1}$.

Proof. This question is equivalent to ask: How many ways are there of distributing k sweets to n children such that each child has at least one sweet.

Lay out the sweets in a single row of length k , and cut it into n pieces. Then give the sweets of the i th piece to child i , which means that we need $n - 1$ cuts from $k - 1$ possibilities. ■

Fact 1.5. The number of integer solutions (x_1, \dots, x_n) to the equation $x_1 + \dots + x_n = k$ with each $x_i \geq 0$ is $\binom{n+k-1}{n-1}$.

Proof 1. Let $A = \{\text{integer solutions } (x_1, \dots, x_n) \text{ to } x_1 + \dots + x_n = k, x_i \geq 0\}$ and $B = \{\text{integer solutions } (y_1, \dots, y_n) \text{ to } y_1 + \dots + y_n = n + k, y_i > 0\}$. Then $|B| = \binom{n+k-1}{n-1}$ by Fact 1.4.

Define $f : A \rightarrow B$, by $f((x_1, \dots, x_n)) = (x_1 + 1, \dots, x_n + 1)$. It suffices to check that f is a bijection, which we omit here. ■

Proof 2. Suppose we have k sweets (of the same sort), which we want to distribute to n children. In how many ways can we do this? Let x_i denote the number of sweets we give to the i -th child, this question is equivalent to that state above.

We lay out the sweets in a single row of length r and let the first child pick them up from left to right (can be 0). After a while we stop him/her and let the second child pick up sweets, etc. The distribution is determined by the specifying the place of where to start a new child. This is equal to select $n - 1$ elements from $n + r - 1$ elements to be the child, others be the sweets (the first child always starts at the beginning). So the answer is $\binom{n+k-1}{n-1}$. ■

Exercise 1.6. Let $X = [n]$, $A = \{(a_1, a_2, \dots, a_r) \mid a_i \in X, 1 \leq a_1 \leq a_2 \leq \dots \leq a_r \leq n, a_{i+1} - a_i \geq k + 1, i \in [r - 1]\}$. Prove that $|A| = \binom{n-k(r-1)}{r}$.

Exercise 1.7. Give a Combinatorial proof of

$$\sum_{k=0}^n \binom{n}{k} \binom{k}{m} = \binom{n}{m} 2^{n-m}.$$

Exercise 1.8. Give a Combinatorial proof of

$$\sum_{k=0}^m \binom{m}{k} \binom{n+k}{m} = \sum_{k=0}^m \binom{n}{k} \binom{m}{k} 2^k.$$

1.2 Counting Mappings

Define X^Y to be the set of all functions $f : Y \rightarrow X$.

Fact 1.9. $|X^Y| = |X|^{|Y|}$.

Proof. Let $|Y| = r$. We can view X^Y as the set of all strings $x_1 x_2 \dots x_r$ with elements $x_i \in X$, indexed by the r elements of Y . So $|X^Y| = |X|^{|Y|}$. ■

Fact 1.10. The number of injective functions $f : [r] \rightarrow [n]$ is $(n)_r$.

Proof. We can view the injective function f as an ordered k -tuple (x_1, x_2, \dots, x_r) with distinct $x_i \in X$, so the number of injective functions $f : [r] \rightarrow [n]$ is $(n)_r$. ■

Definition 1.11 (The Stirling number of the second kind). Let $S(r, n)$ be the number of partitions of $[r]$ into n unordered non-empty parts.

Exercise 1.12. Prove that

$$S(r, 2) = \frac{2^r - 2}{2} = \frac{1}{2} \sum_{i=1}^{r-1} \binom{r}{i}.$$

Fact 1.13. The number of surjective functions $f : [r] \rightarrow [n]$ is $n!S(r, n)$.

Proof. Since f is a surjective function if and only if for any $i \in [n]$, $f^{-1}(i) \neq \emptyset$ if and only if $\cup_{i \in [n]} f^{-1}(i) = [r]$, and $S(r, n)$ is the number of partition of $[r]$ into n unordered non-empty parts, we have the number of surjective functions $f : [r] \rightarrow [n]$ is $n!S(r, n)$. ■

We say that any injective $f : X \rightarrow X$ is a **permutation** of X (also a bijection). We may view a permutation in two ways: (1) it is a bijective from X to X . (2) a reordering of X .

Cycle notation describes the effect of repeatedly applying the permutation on the elements of the set. It expresses the permutation as a product of cycles; since distinct cycles are disjoint, this is referred to as “decomposition into disjoint cycles”.

Definition 1.14 (The Stirling number of the first kind). Let $s(r, n)$ be the number of permutations of $[r]$ with exactly n cycles multiplied by $(-1)^{(r-n)}$.

The following fact is a direct consequence of Fact 1.10.

Fact 1.15. *The number of permutations of $[n]$ is $n!$.*

Exercise 1.16. (1) Let $S(r, n) = \left\{ \begin{matrix} r \\ n \end{matrix} \right\}$, give a Combinatorial proof of $\left\{ \begin{matrix} n \\ k \end{matrix} \right\} = \left\{ \begin{matrix} n-1 \\ k-1 \end{matrix} \right\} + k \left\{ \begin{matrix} n-1 \\ k \end{matrix} \right\}$.

(2) Let $s(n, k) = (-1)^{n-k} \left[\begin{matrix} n \\ k \end{matrix} \right]$, give a Combinatorial proof of $\left[\begin{matrix} n \\ k \end{matrix} \right] = \left[\begin{matrix} n-1 \\ k-1 \end{matrix} \right] + (n-1) \left[\begin{matrix} n-1 \\ k \end{matrix} \right]$.

1.3 The Binomial Theorem

Define $[x^k]f$ to be the coefficient of the term x^k in the polynomial $f(x)$.

Fact 1.17. *For $j = 1, 2, \dots, n$, let $f_j(x) = \sum_{k \in I_j} x^k$ where I_j is a set of non-negative integers, and let $f(x) = \prod_{j=1}^n f_j(x)$. Then, $[x^k]f$ equals the number of solutions (i_1, i_2, \dots, i_n) to $i_1 + i_2 + \dots + i_n = k$, where $i_j \in I_j$.*

Fact 1.18. *Let f_1, \dots, f_n be polynomials and $f = f_1 f_2 \dots f_n$. Then,*

$$[x^k]f = \sum_{i_1 + \dots + i_n = k, i_j \geq 0} \left(\prod_{j=1}^n [x^{i_j}]f_j \right).$$

Theorem 1.19 (The Binomial Theorem). *For any real x and any positive integer n , we have*

$$(1+x)^n = \sum_{i=0}^n \binom{n}{i} x^i.$$

Proof 1. Let $f = (1+x)^n$. By Fact 1.17 we have $[x^k]f$ equals the number of solutions (i_1, i_2, \dots, i_n) to $i_1 + i_2 + \dots + i_n = k$ where $i_j \in \{0, 1\}$, so $[x^k]f = \binom{n}{k}$. ■

Proof 2. By induction on n . When $n = 1$, it is trivial. If the result holds for $n - 1$, then $(1+x)^n = (1+x)(1+x)^{n-1} = (1+x) \sum_{i=0}^{n-1} \binom{n-1}{i} x^i = \sum_{i=1}^n \left(\binom{n-1}{i} + \binom{n-1}{i-1} \right) x^i + 1 + x^n$. Since $\binom{n-1}{i} + \binom{n-1}{i-1} = \binom{n}{i}$ and $\binom{n}{0} = \binom{n}{n} = 1$, we have $(1+x)^n = \sum_{i=0}^n \binom{n}{i} x^i$. ■

Fact 1.20. $\binom{2n}{n} = \sum_{i=0}^n \binom{n}{i}^2 = \sum_{i=0}^n \binom{n}{i} \binom{n}{n-i}$.

Proof 1. Since $(1+x)^{2n} = (1+x)^n (1+x)^n$, by Fact 1.18, we have $\binom{2n}{n} = [x^n](1+x)^{2n} = \sum_{i=0}^n ([x^i](1+x)^n) ([x^{n-i}](1+x)^n) = \sum_{i=0}^n \binom{n}{i} \binom{n}{n-i} = \sum_{i=0}^n \binom{n}{i}^2$. ■

Proof 2. (It is easy to find a combinatorial proof.) ■

Exercise 1.21 (Vandermonde's Convolution Formula).

$$\binom{n+m}{k} = \sum_{j=0}^k \binom{n}{j} \binom{m}{k-j} = \sum_{i+j=k} \binom{n}{i} \binom{m}{j}.$$

Exercise 1.22.

$$\binom{n+m}{r+m} = \sum_{i-j=r} \binom{n}{i} \binom{m}{j}.$$

Exercise 1.23. Prove that

$$\sum_{k=0}^m \binom{m}{k} \binom{n+k}{m} = \sum_{k=0}^m \binom{n}{k} \binom{m}{k} 2^k.$$

by Binomial Theorem.

Fact 1.24. (1).

$$\sum_{\text{all even } k} \binom{n}{k} = \sum_{\text{all odd } k} \binom{n}{k} = 2^{n-1}.$$

(2).

$$\sum_{k=0}^n k \binom{n}{k} = n2^{n-1}.$$

Proof. (1). We see that $(1+x)^n = \sum_{i=0}^n \binom{n}{i} x^i$. Taking $x = 1$ and $x = -1$, we have

$$\sum_{\text{all even } k} \binom{n}{k} = \sum_{\text{all odd } k} \binom{n}{k} = 2^{n-1}.$$

(2). Let $f(x) = (1+x)^n = \sum_{k=0}^n \binom{n}{k} x^k$. Then $f'(x) = n(1+x)^{n-1} = \sum_{k=0}^n k \binom{n}{k} x^{k-1}$. Let $x = 1$, then we have $\sum_{k=0}^n k \binom{n}{k} = n2^{n-1}$. ■

Definition 1.25. Let $k_j \geq 0$ be integers satisfying that $k_1 + k_2 + \dots + k_m = n$. We define

$$\binom{n}{k_1, k_2, \dots, k_m} := \frac{n!}{k_1! k_2! \dots k_m!}.$$

- When $m = 2$, $\binom{n}{k_1, k_2} = \binom{n}{k_1}$ is the number of binary vectors of length n with k_1 zero and k_2 ones, which is also the number of ordered partitions of $[n]$ into 2 parts such that the i_{th} part has size k_i .
- When $m \geq 3$, $\binom{n}{k_1, k_2, \dots, k_m}$ is the number of m -ary vectors of length n over $[m]$ such that i occurs k_i times, which is also the number of ordered partitions of $[n]$ into m parts such that the i_{th} part has size k_i .

The following theorem is a generalization of the binomial theorem.

Exercise 1.26 (Multinomial Theorem). For any reals x_1, \dots, x_m and any positive integer $n \geq 1$, we have

$$(x_1 + x_2 + \dots + x_m)^n = \sum_{k_1+k_2+\dots+k_m=n, k_j \geq 0} \binom{n}{k_1, k_2, \dots, k_m} x_1^{k_1} x_2^{k_2} \dots x_m^{k_m}.$$

Exercise 1.27. Suppose $\sum_{i=1}^m k_i = n$ with $k_i \geq 1$ for all $i \in [m]$. Then

$$\binom{n}{k_1, k_2, \dots, k_m} = \binom{n-1}{k_1-1, k_2, \dots, k_m} + \dots + \binom{n-1}{k_1, k_2, \dots, k_m-1}.$$

1.4 Inclusion and Exclusion Principle (IEP)

This lecture is devoted to Inclusion-Exclusion formula and its applications.

Let Ω be a ground set and let A_1, A_2, \dots, A_n be subsets of Ω . Write $A_i^c = \Omega \setminus A_i$. Throughout this lecture, we use the following notation.

Definition 1.28. Let $A_\emptyset = \Omega$. For any nonempty subset $I \subseteq [n]$, let

$$A_I = \bigcap_{i \in I} A_i.$$

For any integer $k \geq 0$, let

$$S_k = \sum_{I \in \binom{[n]}{k}} |A_I|.$$

Now we introduce Inclusion-Exclusion formula (in three equivalent forms) and give two proofs as follows.

Theorem 1.29 (Inclusion-Exclusion Formula). *We have*

$$|A_1 \cup A_2 \cup \dots \cup A_n| = \sum_{k=1}^n (-1)^{k+1} S_k,$$

which is equivalent to

$$\left| \Omega \setminus \bigcup_{i=1}^n A_i \right| = |A_1^c \cap A_2^c \cap \dots \cap A_n^c| = \sum_{k=0}^n (-1)^k S_k,$$

and

$$\left| \Omega \setminus \bigcup_{i=1}^n A_i \right| = |A_1^c \cap A_2^c \cap \dots \cap A_n^c| = \sum_{I \subseteq [n]} (-1)^{|I|} |A_I|.$$

Proof (1). For any subset $X \subseteq \Omega$, we define its characterization function $\mathbb{1}_X : \Omega \rightarrow \{0, 1\}$ by assigning

$$\mathbb{1}_X(x) = \begin{cases} 1, & x \in X \\ 0, & x \notin X. \end{cases}$$

Then we notice that $\sum_{x \in \Omega} \mathbb{1}_X(x) = |X|$. Let $A = A_1 \cup A_2 \cup \dots \cup A_n$. Our key observation is that

$$(\mathbb{1}_A - \mathbb{1}_{A_1})(\mathbb{1}_A - \mathbb{1}_{A_2}) \cdots (\mathbb{1}_A - \mathbb{1}_{A_n})(x) \equiv 0,$$

which holds for any $x \in \Omega$. Next we expand this product into a summation of 2^n terms as follows:

$$\mathbb{1}_A + \sum_{\emptyset \neq I \subseteq [n]} (-1)^{|I|} \left(\prod_{i \in I} \mathbb{1}_{A_i} \right) \equiv 0$$

holds for any $x \in \Omega$. Summing over all $x \in \Omega$, this gives that

$$|A| + \sum_{\emptyset \neq I \subseteq [n]} (-1)^{|I|} |A_I| = 0,$$

which implies that

$$|A_1 \cup A_2 \cup \dots \cup A_n| = |A| = \sum_{\emptyset \neq I \subseteq [n]} (-1)^{|I|+1} |A_I| = \sum_{k=1}^n (-1)^{k+1} S_k,$$

finishing the proof. ■

Proof (2). It suffices to prove that

$$\mathbb{1}_{A_1 \cup A_2 \cup \dots \cup A_n}(x) = \sum_{k=1}^n (-1)^{k+1} \sum_{I \in \binom{[n]}{k}} \mathbb{1}_{A_I}(x)$$

holds for all $x \in \Omega$. Denote by LHS (resp. RHS) the left-hand side (resp. right-hand side) of the above equation.

Assume that x is contained in exactly ℓ subsets, say A_1, A_2, \dots, A_ℓ . If $\ell = 0$, then clearly $LHS = 0 = RHS$, so we are done. So we may assume that $\ell \geq 1$. In this case, we have $LHS = 1$ and

$$RHS = \ell - \binom{\ell}{2} + \binom{\ell}{3} + \dots + (-1)^{\ell+1} \binom{\ell}{\ell} = 1.$$

Note that the above equation holds since $\sum_{i=0}^{\ell} (-1)^i \binom{\ell}{i} = (1-1)^\ell = 0$. This finishes the proof. ■

Next, we will demonstrate the power of Inclusion-Exclusion formula by using it to solve several problems.

Definition 1.30. Let $\varphi(n)$ be the number of integers $m \in [n]$ which are relatively prime¹ to n .

Theorem 1.31. If we express $n = p_1^{a_1} p_2^{a_2} \dots p_t^{a_t}$, where p_1, \dots, p_t are distinct primes, then

$$\varphi(n) = n \prod_{i=1}^t \left(1 - \frac{1}{p_i}\right).$$

Proof. Let the ground set

$$\Omega = [n]$$

and

$$A_i = \{m \in [n] : p_i | m\}$$

for $i \in \{1, 2, \dots, t\}$. It implies

$$\varphi(n) = |\{m \in [n] : m \notin A_i \text{ for all } i \in [t]\}| = |[n] \setminus (A_1 \cup A_2 \cup \dots \cup A_t)|.$$

By Inclusion-Exclusion formula,

$$\varphi(n) = \sum_{I \subseteq [t]} (-1)^{|I|} |A_I|,$$

¹Here, “ m is relatively prime to n ” means that the greatest common divisor of m and n is 1.

where $A_I = \cap_{i \in I} A_i = \{m \in [n] : (\prod_{i \in I} p_i) | m\}$ and thus $|A_I| = \frac{n}{\prod_{i \in I} p_i}$. We can derive that

$$\varphi(n) = \sum_{I \subseteq [t]} (-1)^{|I|} \frac{n}{\prod_{i \in I} p_i} = n \left(1 - \frac{1}{p_1}\right) \left(1 - \frac{1}{p_2}\right) \cdots \left(1 - \frac{1}{p_t}\right),$$

as desired. ■

Exercise 1.32. For any positive integer n ,

$$\sum_{d|n} \varphi(d) = n.$$

1.5 Möbius Inversion Formula

Definition 1.33. The Möbius Function μ for a positive integer d is

$$\mu(d) = \begin{cases} 1, & d \text{ is a product of even number of distinct primes } (d = 1 \text{ included}) \\ -1, & d \text{ is a product of odd number of distinct primes} \\ 0, & \text{otherwise} \end{cases}$$

Theorem 1.34. For any positive integer n ,

$$\sum_{d|n} \mu(d) = \begin{cases} 1, & n = 1 \\ 0, & \text{otherwise} \end{cases}$$

Proof. If $n = 1$, it is trivial. For $n = p_1^{a_1} \cdots p_r^{a_r} \geq 2$,

$$\sum_{d|n} \mu(d) = \sum_{i_1 \leq a_1, \dots, i_r \leq a_r} \mu(p_1^{i_1} \cdots p_r^{i_r}) = \sum_{i=0}^r \binom{r}{i} (-1)^i = 0.$$

■

Theorem 1.35 (Möbius Inversion Formula). Let $f(n)$ and $g(n)$ be two functions defined for every positive integer n satisfying

$$f(n) = \sum_{d|n} g(d).$$

Then we have

$$g(n) = \sum_{d|n} \mu(d) f\left(\frac{n}{d}\right).$$

Proof.

$$\begin{aligned}
\sum_{d|n} \mu(d) f\left(\frac{n}{d}\right) &= \sum_{d|n} \mu\left(\frac{n}{d}\right) f(d) \\
&= \sum_{d|n} \mu\left(\frac{n}{d}\right) \left(\sum_{d'|d} g(d')\right) \\
&= \sum_{d'|n} g(d') \sum_{\frac{n}{d} | \frac{n}{d'}} \mu\left(\frac{n}{d}\right) \\
&= \sum_{d'|n} g(d') \sum_{m | \frac{n}{d'}} \mu(m) \\
&= \sum_{d'|n, d' \neq n} g(d') \times 0 + g(n) \times 1 \\
&= g(n)
\end{aligned}$$

as desired. ■

1.6 Generating Functions

Definition 1.36. *The (ordinary) generating function (GF) for an infinite sequence $\{a_0, a_1, \dots\}$ is a power series*

$$f(x) = \sum_{n \geq 0} a_n x^n.$$

We have two ways to view this power series.

- (i). When the power series $\sum_{n \geq 0} a_n x^n$ converges (i.e. there exists a radius $R > 0$ of convergence), we view GF as a function of x and we can apply operations of calculus on it (including derivation and integration). For example, we know that

$$a_n = \frac{f^{(n)}(0)}{n!}.$$

Recall the following sufficient condition on the radius of convergence that if $|a_n| \leq K^n$ for some $K > 0$, then $\sum_{n \geq 0} a_n x^n$ converges in the interval $(-\frac{1}{K}, \frac{1}{K})$.

- (ii). When we are not sure of the convergence, we view the generating function as a formal series and take additions and multiplications. Let $a(x) = \sum_{n \geq 0} a_n x^n$ and $b(x) = \sum_{n \geq 0} b_n x^n$.

Addition.

$$a(x) + b(x) = \sum_{n \geq 0} (a_n + b_n) x^n.$$

Multiplication. Let $c_n = \sum_{i=0}^n a_i b_{n-i}$. Then

$$a(x) \cdot b(x) = \sum_{n \geq 0} c_n x^n.$$

Example 1.37. *Consider the GF of $\{1, 1, 1, \dots\}$. We note $\frac{1}{1-x} = \sum_{n=0}^{\infty} x^n$ holds for all $-1 < x < 1$. From the point view of (i), its first derivative gives*

$$\frac{1}{(1-x)^2} = \sum_{n=1}^{\infty} n x^{n-1} = \sum_{n=0}^{\infty} (n+1) x^n.$$

So we could view $\frac{1}{(1-x)^2}$ as the GF of $\{1, 2, 3, \dots\}$ for all $-1 < x < 1$.

Problem 1.38. Let $a_0 = 1$ and $a_n = 2a_{n-1}$ for $n \geq 1$. Find a_n .

Solution. Consider the generating function,

$$f(x) = \sum_{n=0}^{\infty} a_n x^n = 1 + \sum_{n=1}^{\infty} a_n x^n = 1 + 2x \sum_{n=1}^{\infty} a_{n-1} x^{n-1} = 1 + 2xf(x).$$

So $f(x) = \frac{1}{1-2x}$, which implies that $f(x) = \sum_{n=0}^{+\infty} 2^n x^n$ and $a_n = 2^n$. ■

From this problem, we see one of the basic ideas for using generating function: in order to find the general expression of a_n , we work on its generating function $f(x)$; once we find the formula of $f(x)$, then we can expand $f(x)$ into a power series and get a_n by choosing the coefficient of the right term.

Problem 1.39. Let A_n be the set of strings of length n with entries from the set $\{a, b, c\}$ and with no “aa” occurring (in the consecutive positions). Find $|A_n|$ for $n \geq 1$.

Solution. Let $a_n = |A_n|$. We first observe that $a_1 = 3, a_2 = 8$. For $n \geq 3$, we will find a_n by recursion as follows. If the first string is ‘a’, the second string has two choices, ‘b’ or ‘c’. Then the last $n - 2$ strings have a_{n-2} choices. If the first string is ‘b’ or ‘c’, the last $n - 1$ strings have a_{n-1} choices. They are all different. Totally, for $n \geq 3$, we have

$$a_n = 2a_{n-1} + 2a_{n-2}.$$

Set $a_0 = 1$, then $a_n = 2a_{n-1} + 2a_{n-2}$ holds for $n \geq 2$. The generating function of $\{a_n\}$ is

$$f(x) = \sum_{n \geq 0} a_n x^n = a_0 + a_1 x + \sum_{n \geq 2} (2a_{n-1} + 2a_{n-2}) x^n = 1 + 3x + 2x(f(x) - 1) + 2x^2 f(x),$$

which implies that

$$f(x) = \frac{1+x}{1-2x-2x^2}.$$

By Partial Fraction Decomposition, we calculate that

$$f(x) = \frac{1-\sqrt{3}}{2\sqrt{3}} \frac{1}{\sqrt{3}+1+2x} + \frac{1+\sqrt{3}}{2\sqrt{3}} \frac{1}{\sqrt{3}-1-2x},$$

which implies that

$$a_n = \frac{1-\sqrt{3}}{2\sqrt{3}} \frac{1}{\sqrt{3}+1} \left(\frac{-2}{\sqrt{3}+1} \right)^n + \frac{1+\sqrt{3}}{2\sqrt{3}} \frac{1}{\sqrt{3}-1} \left(\frac{2}{\sqrt{3}-1} \right)^n.$$

■

Remark 1.40. Note that a_n must be an integer but its expression is a combination of irrational terms! Observe that $\left| \frac{-2}{\sqrt{3}+1} \right| < 1$, so $\left(\frac{-2}{\sqrt{3}+1} \right)^n \rightarrow 0$ as $n \rightarrow \infty$. Thus, when n is sufficiently large, this integer a_n is about the value of the second term $\frac{1+\sqrt{3}}{2\sqrt{3}} \frac{1}{\sqrt{3}-1} \left(\frac{2}{\sqrt{3}-1} \right)^n$. Equivalently a_n will be the nearest integer to that.

Exercise 1.41. Define Fibonacci number F_n as follows: $F_1 = 0, F_2 = 1, F_{n+2} = F_{n+1} + F_n$ for all $n \geq 0$. Find F_n .

Definition 1.42. For any real r and an integer $k \geq 0$, let

$$\binom{r}{k} = \frac{r(r-1)\cdots(r-k+1)}{k!}.$$

Exercise 1.43. Prove that $\binom{\frac{1}{2}}{k} = \frac{(-1)^{k-1} \cdot 2 \cdot (2k-2)!}{4^k \cdot k!(k-1)!}$.

Theorem 1.44 (Newton's Binomial Theorem). For any real number r and $x \in (-1, 1)$,

$$(1+x)^r = \sum_{k=0}^{\infty} \binom{r}{k} x^k.$$

Proof. By Taylor series, it is obvious. ■

Corollary 1.45. Let $r = -n$ for some integer $n \geq 0$. Then

$$\binom{-n}{k} = \frac{(-n)(-n-1)\cdots(-n-k+1)}{k!} = (-1)^k \binom{n+k-1}{k}.$$

Therefore

$$(1+x)^{-n} = \sum_{k=0}^{\infty} (-1)^k \binom{n+k-1}{k} x^k,$$

which is equivalent to

$$(1-x)^{-n} = \sum_{k=0}^{\infty} \binom{n+k-1}{k} x^k.$$

Noting that

$$\binom{n+k-1}{k} = \# \text{ integer solutions to } x_1 + x_2 + \cdots + x_n = k \text{ where } x_i \geq 0, 1 \leq i \leq n,$$

we can explain Equation (3.21) from another point of view as follows.

Recall the following facts.

Fact 1.46. For $j \in [n]$, let $f_j(x) := \sum_{i \in I_j} x^i$, where $I_j \subset \mathbb{N}$. Let b_k be the number of solutions to $i_1 + i_2 + \cdots + i_n = k$ for $i_j \in I_j$. Then

$$\prod_{j=1}^n f_j(x) = \sum_{k=0}^{\infty} b_k x^k.$$

Fact 1.47. If $f(x) = \prod_{i=1}^k f_i(x)$ for polynomials f_1, \dots, f_k , then

$$[x^n]f = \sum_{i_1+i_2+\cdots+i_k=n} \prod_{j=1}^k ([x^{i_j}]f_j),$$

where $[x^n]f$ is the coefficient of x^n in f .

Let $f_j = (1-x)^{-1} = \sum_{i \geq 0} x^i, \forall j \in [n]$. By Fact 1.46, we can get Equation 3.21 by considering as $(1-x)^{-n} = \prod_{j=1}^n f_j$ easily.

Exercise 1.48. Show $(1-x)^{-n} = \sum_{k=0}^{\infty} \binom{n+k-1}{k} x^k$ by taking the n^{th} derivative of $(1-x)^{-1}$.

Problem 1.49. Let a_n be the number of ways to pay n Yuan using 1-Yuan bills, 2-Yuan bills and 5-Yuan bills. What is the generating function of this sequence $\{a_n\}$?

Solution. Observe that a_n is the number of integer solutions (i_1, i_2, i_3) to $i_1 + i_2 + i_3 = n$, where $i_1 \in I_1 := \{0, 1, 2, \dots\}$, $i_2 \in I_2 := \{0, 2, 4, \dots\}$ and $i_3 \in I_3 := \{0, 5, 10, \dots\}$. Let $f_j(x) := \sum_{m \in I_j} x^m$ for $j = 1, 2, 3$. By Fact 1.46, we have

$$\sum_{n=0}^{+\infty} a_n x^n = f_1(x) f_2(x) f_3(x) = \frac{1}{1-x} \cdot \frac{1}{1-x^2} \cdot \frac{1}{1-x^5}.$$

■

1.7 Random Walks

Consider a real axis with integer points $(0, \pm 1, \pm 2, \pm 3, \dots)$ marked. A frog leaps among the integer points according to the following rules:

- (1). At beginning, it sits at 1.
- (2). In each coming step, the frog leaps either by distance 2 to the right (from i to $i+2$), or by distance 1 to the left (from i to $i-1$), each of which is randomly chosen with probability $\frac{1}{2}$ independently of each other.

Problem 1.50. What is the probability that the frog can reach “0”?

Solution. In each step, we use “+” or “-” to indicate the choice of the frog that is either to leap right or leap left. Then the probability space Ω can be viewed as the set of infinite vectors, where each entry is in $\{+, -\}$.

Let A be the event that the frog reaches “0”. Let A_i be the event that the frog reaches “0” at the i^{th} step for the first time. So $A = \cup_{i=1}^{+\infty} A_i$ is a disjoint union. So $P(A) = \sum_{i=1}^{+\infty} P(A_i)$.

To compute $P(A_i)$, we can define a_i to be the number of trajectories (or vectors) of the first i steps such that the frog starts at “1” and reaches “0” at the i^{th} step for the first time. So

$$P(A_i) = \frac{a_i}{2^i}.$$

Then,

$$P(A) = \sum_{i=1}^{+\infty} \frac{a_i}{2^i}.$$

Let $f(x) = \sum_{i=0}^{+\infty} a_i x^i$ be the generating function of $\{a_i\}_{i \geq 0}$, where $a_0 := 0$. Thus,

$$P(A) = \sum_{i=1}^{+\infty} \frac{a_i}{2^i} = f\left(\frac{1}{2}\right).$$

We then turn to find the expression of $f(x)$.

Let b_i be the number of trajectories of the first i steps such that the frog starts at “2” and reaches “0” at the i^{th} step for the first time.

Let c_i be the number of trajectories of the first i steps such that the frog starts at “3” and reaches “0” at the i^{th} step for the first time.

First we express b_i in terms of $\{a_j\}_{j \geq 1}$. Since the frog only can leap to left by distance 1, if the frog can successfully jump from “ i ” to “0” in i steps, then this frog must reach “1” first. Let j be the number of steps by which the frog reaches “1” for the first time. So there are a_j trajectories from “2” to “1” at the j^{th} step for the first time. In the remaining $i - j$ steps the frog must jump from “1” to “0” and reach “0” at the coming $(i - j)^{\text{th}}$ step for the first time, so there are a_{i-j} trajectories that the frog can finish in exactly $i - j$ steps. In total,

$$b_i = \sum_{j=1}^{i-1} a_j a_{i-j}.$$

As $a_0 = 0$,

$$b_i = \sum_{j=0}^i a_j a_{i-j}.$$

We can get

$$\sum_{i \geq 0} b_i x^i = \left(\sum_{i \geq 0} a_i x^i \right)^2 = f^2(x).$$

Similarly, if we count the number c_i of trajectories from 3 to 0, we can obtain that

$$c_i = \sum_{j=0}^i a_j b_{i-j},$$

which implies that

$$\sum_{i \geq 0} c_i x^i = \left(\sum_{i \geq 0} b_i x^i \right) \left(\sum_{i \geq 0} a_i x^i \right) = f^3(x).$$

Let us consider a_i from another point of view. After the first step, either the frog reaches “0” directly (if it leaps to left, so $a_1 = 1$), or it leaps to “3”. In the latter case, the frog needs to jump from “3” to “0” using $i - 1$ steps. Thus for $i \geq 2$, $a_i = c_{i-1}$.

Combining the above facts, we have

$$f(x) = \sum_{i=0}^{+\infty} a_i x^i = x + \sum_{i \geq 2} a_i x^i = x + \sum_{i \geq 2} c_{i-1} x^i = x + x \left(\sum_{j=0}^{+\infty} c_j x^j \right) = x + x \cdot f^3(x).$$

Let $a := P(A) = f(1/2)$. Then we have $a = \frac{1}{2} + \frac{a^3}{2}$, i.e., $(a - 1)(a^2 + a - 1) = 0$, implying that

$$a = 1, \frac{\sqrt{5} - 1}{2} \text{ or } \frac{-\sqrt{5} - 1}{1}.$$

Since $P(A) \in [0, 1]$, we see $P(A) = 1$ or $\frac{\sqrt{5}-1}{2}$.

Note that $f(x) = x + xf^3(x)$. Consider the inverse function of $f(x)$, that is, $g(x) := \frac{x}{1+x^3}$. Consider the figure of $g(x)$. We find that $g(x)$ is increasing around $\frac{\sqrt{5}-1}{2}$ but decreasing around 1. Since $f(x) = \sum a_i x^i$ is increasing, $g(x)$ also increases. Thus it doesn't make sense for $g(x)$ being around $x = 1$. This explains that $P(A) = \frac{\sqrt{5}-1}{2}$, which is the golden section! ■

1.8 Exponential Generating Functions

Let \mathbb{N} , \mathbb{N}_e and \mathbb{N}_o be the sets of non-negative integers, non-negative even integers and non-negative odd integers, respectively.

Given n sets I_j of non-negative integers for $j \in [n]$, let $f_j(x) = \sum_{i \in I_j} x^i$. Let a_k be the number of integer solutions to $i_1 + i_2 + \dots + i_n = k$, where $i_j \in I_j$. Then $\prod_{j=1}^n f_j(x)$ is the ordinary generating function of $\{a_k\}_{k \geq 0}$.

Problem 1.51. Let S_n be the number of selections of n letters chosen from an unlimited supply of a 's, b 's and c 's such that both of the numbers of a 's and b 's are even.

Solution. We can write S_n as

$$S_n = \sum_{e_1+e_2+e_3=n, e_1, e_2 \in \mathbb{N}_e, e_3 \in \mathbb{N}} 1.$$

Using the previous fact, we see that $S_n = [x^n]f$, where

$$f(x) = \left(\sum_{i \in \mathbb{N}_e} x^i \right)^2 \left(\sum_{j \in \mathbb{N}} x^j \right) = \left(\frac{1}{1-x^2} \right)^2 \cdot \frac{1}{1-x}.$$

■

Problem 1.52. Let T_n be the number of arrangements (or words) of n letters chosen from an unlimited supply of a 's, b 's and c 's such that both of the numbers of a 's and b 's are even. What is the value of T_n ?

Solution. To solve this, we define a new kind of generating functions.

Definition 1.53. The exponential generating function for the sequence $\{a_n\}_{n \geq 0}$ is the power series

$$f(x) = \sum_{n=0}^{\infty} a_n \cdot \frac{x^n}{n!}.$$

Then we have the following fact.

Fact 1.54. If we have n letters including x a 's, y b 's and z c 's (i.e. $x + y + z = n$), then we can form $\frac{n!}{x!y!z!}$ distinct words using them.

Therefore, a selection (say x a 's, y b 's and z c 's) can contribute $\frac{n!}{x!y!z!}$ arrangements to T_n . This implies that

$$T_n = \sum_{e_1+e_2+e_3=n, e_1, e_2 \in \mathbb{N}_e, e_3 \in \mathbb{N}} \frac{n!}{e_1!e_2!e_3!}.$$

Similar to defining the above $f(x)$ for S_n , we define the following for T_n . Let

$$g(x) := \left(\sum_{i \in \mathbb{N}_e} \frac{x^i}{i!} \right)^2 \left(\sum_{j \in \mathbb{N}} \frac{x^j}{j!} \right).$$

Claim. We have

$$[x^n]g = \frac{T_n}{n!}.$$

Proof. To see this, we expand $g(x)$. Then the term x^n in $g(x)$ becomes

$$\sum_{\substack{e_1+e_2+e_3=n, \\ e_1, e_2 \in \mathbb{N}_e, e_3 \in \mathbb{N}}} \frac{x^{e_1}}{e_1!} \cdot \frac{x^{e_2}}{e_2!} \cdot \frac{x^{e_3}}{e_3!} = \left(\sum_{\substack{e_1+e_2+e_3=n, \\ e_1, e_2 \in \mathbb{N}_e, e_3 \in \mathbb{N}}} \frac{n!}{e_1!e_2!e_3!} \right) \frac{x^n}{n!} = T_n \cdot \frac{x^n}{n!}.$$

So $[x^n]g = \frac{T_n}{n!}$, i.e., $g(x)$ is the exponential generating function of $\{T_n\}$. This finishes the proof of Claim. \blacksquare

Using Taylor series: $e^x = \sum_{j \geq 0} \frac{x^j}{j!}$ and $e^{-x} = \sum_{j \geq 0} (-1)^j \frac{x^j}{j!}$, we have

$$\frac{e^x + e^{-x}}{2} = \sum_{j \in \mathbb{N}_e} \frac{x^j}{j!} \quad \text{and} \quad \frac{e^x - e^{-x}}{2} = \sum_{j \in \mathbb{N}_o} \frac{x^j}{j!}.$$

By the previous fact, we get

$$g(x) = \left(\frac{e^x + e^{-x}}{2} \right)^2 \cdot e^x = \frac{e^{3x} + 2e^x + e^{-x}}{4} = \sum_{n \geq 0} \left(\frac{3^n + 2 + (-1)^n}{4} \right) \cdot \frac{x^n}{n!}.$$

Therefore, we get that

$$T_n = \frac{3^n + 2 + (-1)^n}{4}.$$

Recall that the *exponential generating function* for the sequence $\{a_n\}_{n \geq 0}$ is the power series

$$f(x) = \sum_{n=0}^{+\infty} a_n \cdot \frac{x^n}{n!}.$$

As we shall see, ordinary generation functions can be used to find the number of selections; while exponential generation functions can be used to find the number of arrangements or some combinatorial objects **involving ordering**. We summarize this as the following facts.

Fact 1.55. Given $I_j \subseteq \mathbb{N}$ for $j \in [n]$, let $f_j(x) = \sum_{i \in I_j} x^i$. And let $a_k = \sum_{\substack{i_1 + \dots + i_n = k, \\ i_j \in I_j}} 1$. Then

$$\prod_{j=1}^n f_j(x) = \sum_{k=0}^{+\infty} a_k x^k.$$

Fact 1.56. Given $I_j \subseteq \mathbb{N}$ for $j \in [n]$, let $g_j(x) = \sum_{i \in I_j} \frac{x^i}{i!}$. And let $b_k = \sum_{\substack{i_1 + \dots + i_n = k, \\ i_j \in I_j}} \frac{k!}{i_1! i_2! \dots i_n!}$. Then

$$\prod_{j=1}^n g_j(x) = \sum_{k=0}^{+\infty} \frac{b_k}{k!} x^k.$$

Fact 1.57. Let $f(x) = \prod_{j=1}^n f_j(x)$. Then

$$[x^k]f = \sum_{\substack{i_1 + \dots + i_n = k, \\ i_j \geq 0}} \prod_{j=1}^n [x^{i_j}]f_j.$$

Fact 1.58. Let $f(x) = \prod_{j=1}^n f_j(x)$ and let $f_j(x) = \sum_{k=0}^{+\infty} \frac{a_k^{(j)}}{k!} x^k$. Then

$$f(x) = \sum_{k=0}^{+\infty} \frac{A_k}{k!} x^k,$$

if and only if

$$A_k = \sum_{\substack{i_1 + \dots + i_n = k, \\ i_j \geq 0}} \frac{k!}{i_1! i_2! \dots i_n!} \left(\prod_{j=1}^n a_{i_j}^{(j)} \right).$$

Exercise 1.59. Find the number a_n of ways to send n students to four different classes (say R_1, R_2, R_3, R_4) such that each class has at least one student.

Solution.

$$a_n = \sum_{\substack{i_1 + i_2 + i_3 + i_4 = n, \\ i_j \geq 1}} \frac{n!}{i_1! i_2! i_3! i_4!}.$$

Let $I_j \subseteq \mathbb{N}$ for $j \in [4]$ and $g_j(x) = \sum_{i \geq 1} \frac{x^i}{i!} = e^x - 1$. By Fact 1.56, we have that

$$\sum_{n=0}^{+\infty} \frac{a_n}{n!} x^n = g_1 g_2 g_3 g_4 = \left(\sum_{i \geq 1} \frac{x^i}{i!} \right)^4 = (e^x - 1)^4 = e^{4x} - 4e^{3x} + 6e^{2x} - 4e^x + 1 = \sum_{n=0}^{+\infty} (4^n - 4 \cdot 3^n + 6 \cdot 2^n - 4) \frac{x^n}{n!} + 1.$$

Thus $a_n = 4^n - 4 \cdot 3^n + 6 \cdot 2^n - 4$ for $n \geq 4$. ■

Exercise 1.60. Let a_n be the number of arrangements of type A for a group of n people, and let b_n be the number of arrangements of type B for a group of n people.

Define a new arrangement of n people called type C as follows:

- Divide the n people into 2 groups (say 1st and 2nd).
- Then arrange the 1st group by an arrangement of type A, and arrange the 2nd group by an arrangement of type B.

Let c_n be the number of arrangements of type C of n people. Let $A(x), B(x), C(x)$ be the exponential generation function for $\{a_n\}, \{b_n\}, \{c_n\}$ respectively. Prove that $C(x) = A(x)B(x)$.

Proof. We can easily see that

$$c_n = \sum_{\substack{i+j=n, \\ i,j \geq 0}} \frac{n!}{i!j!} a_i b_j.$$

Then by Fact 1.58, $C(x) = A(x)B(x)$. ■

Exercise 1.61. Recall that $S(n, k) \cdot k!$ is equal to the number of surjections from $[n]$ to $[k]$. For fixed k , compute the exponential generating function of $S(n, k) \cdot k!$. Then find the value of $S(n, k) \cdot k!$.

Fact 1.62 (Lagrange Inversion Formula). Let $f(x)$ be analytic (convergent power series) in a neighborhood of $z = 0$ and $f(0) \neq 0$. If $w = \frac{z}{f(z)}$, then z can be expressed as a power series

$$z = \sum_{k=1}^{\infty} c_k w^k$$

with a positive radius of convergence, where

$$c_k = \frac{1}{k!} \left\{ \left(\frac{d}{dz} \right)^{k-1} (f(z))^k \right\}_{z=0}.$$

2 Basics of Graphs

In this second part of our course, we will introduce some basic definitions about graphs.

Definition 2.1. A graph $G = (V, E)$ consists of a vertex set V and an edge set E , where the elements of V are called **vertices** and the elements of $E \subseteq \binom{V}{2} = \{\{x, y\} : x, y \in V\}$ are called **edges**.

This provides the definition of a simple undirected graph. The word “undirected” means that the edge set E contains unordered pairs. Otherwise, G is called a directed graph. A graph is *simple* if it has no loops or multiple edges. A *loop* is an edge whose endpoints are equal. *Multiple edges* are edges having the same pair of endpoints.

- We say vertices x and y are *adjacent* if $\{x, y\} \in E$, write $x \sim_G y$ or $x \sim y$ or $xy \in E$.
- We say the edge xy is *incident* to the endpoints x and y .
- Let $e(G)$ be the number of edges in G , i.e., $e(G) = |E(G)|$.
- The *degree* of a vertex v in G , denoted by $d_G(v)$, is the number of edges in G incident to v .
- The *neighborhood* of a vertex v is the set of vertices that are adjacent to v , i.e., $N_G(v) = \{u \in V(G) : u \sim v\}$. Thus we have $d_G(v) = |N_G(v)|$.
- A graph $G' = (V', E')$ is a *subgraph* of $G = (V, E)$ if $V' \subseteq V$ and $E' \subseteq E \cap \binom{V'}{2}$, i.e., $G' \subseteq G$.
- A subgraph $G' = (V', E')$ of $G = (V, E)$ is *induced*, if $E' = E \cap \binom{V'}{2}$, write $G' = G[V']$.

Definition 2.2. Two graphs $G = (V, E)$ and $G' = (V', E')$ are **isomorphic** if there exists a bijection $f : V \rightarrow V'$ such that $i \sim_G j$ if and only if $f(i) \sim_{G'} f(j)$.

- A graph on n vertices is a *complete graph* (or a *clique*), denoted by K_n , if all pairs of vertices are adjacent. So we have $e(K_n) = \binom{n}{2}$.
- A graph on n vertices is called an *independent set*, denoted by I_n , if it contains no edge at all.
- Given a graph $G = (V, E)$, its *complement* is a graph $\overline{G} = (V, E^c)$ with $E^c = \binom{V}{2} \setminus E$.
- The *degree sequence* of a graph $G = (V, E)$ is a sequence of degrees of all vertices listed in a non-decreasing order.
- The *path* P_k of length $k - 1$ is a graph $v_1 v_2 \dots v_k$ where $v_i \sim v_{i+1}$ for $i \in [k - 1]$ and $v_j \neq v_l$ for any $j \neq l \in [k]$. Note that the length of a path P (denoted by $|P|$) is the number of edges in P .
- A *cycle* C_k of length k is a graph $v_1 v_2 \dots v_k v_1$ where $v_i \sim v_{i+1}$ for $i \in [k]$, $v_{k+1} = v_1$, and $v_j \neq v_l$ for any $j \neq l \in [k]$.
- Let G be a simple graph with vertex set $V(G) = \{v_1, \dots, v_n\}$ and edge set $E(G) = \{e_1, \dots, e_m\}$. The *adjacency matrix* of G , denoted by $A(G)$, is the n -by- n matrix in which entry $a_{i,j}$ is the number of edges in G with endpoints $\{v_i, v_j\}$. The *incidence matrix* $M(G)$ is the n -by- m matrix in which entry $m_{i,j}$ is 1 if v_j is an endpoint of e_j and 0 otherwise.
- A graph G is *planar*, if we can draw G on the plane such that its edges intersect only at their endpoints.

Theorem 2.3 (Euler's Formula). *Let $G = (V, E)$ be a connected planar graph with v vertices and e edges, and let r be the number of regions in which some given embedding of G divides the plane. Then $v - e + r = 2$.*

Exercise 2.4. *Show that K_4 is planar but K_5 is not.*

Exercise 2.5. *Show that $K_{3,3}$ is not planar.*

The following Handshaking Lemma is the most basic lemma in graph theory.

Lemma 2.6 (Handshaking Lemma). *In any graph $G = (V, E)$,*

$$\sum_{v \in V} d_G(v) = 2e(G).$$

Proof. Let $F = \{(e, v) : e \in E(G), v \in V(G) \text{ such that } v \text{ is incident to } e\}$. Then

$$\sum_{e \in E(G)} 2 = |F| = \sum_{v \in V} d_G(v).$$

■

Corollary 2.7. *In any graph G , the number of vertices with odd degree is even.*

Proof. Let $O = \{v \in V(G) : d(v) \text{ is odd}\}$ and $\mathcal{E} = \{v \in V(G) : d(v) \text{ is even}\}$. Then by Lemma 2.6,

$$2e(G) = \sum_{v \in O} d_G(v) + \sum_{v \in \mathcal{E}} d_G(v).$$

Thus we have $\sum_{v \in O} d_G(v)$ is even, moreover we have $|O|$ is even. ■

Corollary 2.8. *In any graph G , if there exists a vertex with odd degree, then there are at least two vertices with odd degree.*

3 Double-counting Method

3.1 Basics

The basic setting of the double counting technique is as follows. Suppose that we are given two finite sets A and B , and a subset $S \subseteq A \times B$. If $(a, b) \in S$, then we say that a and b are incident. Let N_a be the number of elements $b \in B$ such that $(a, b) \in S$, and N_b be the number of elements $a \in A$ such that $(a, b) \in S$. Then we have

$$\sum_{a \in A} N_a = |S| = \sum_{b \in B} N_b.$$

Theorem 3.1. *Let $T(j)$ be the number of divisions of a positive integer j . Let $\overline{T(n)} = \frac{1}{n} \sum_{j=1}^n T(j)$. Then we have $|\overline{T(n)} - H(n)| < 1$, where $H(n) = \sum_{i=1}^n \frac{1}{i}$ is the n^{th} Harmonic number.*

Proof. Define a table $X = (x_{ij})$ where

$$x_{ij} = \begin{cases} 1, & \text{if } i|j \\ 0, & \text{otherwise.} \end{cases}$$

Then

$$\sum_{j=1}^n T(j) = \sum_{1 \leq i \leq j \leq n} x_{ij} = \sum_{i=1}^n \lfloor \frac{n}{i} \rfloor,$$

which implies that

$$\overline{T(n)} = \frac{1}{n} \sum_{i=1}^n \lfloor \frac{n}{i} \rfloor.$$

Then we have

$$|\overline{T(n)} - H(n)| < 1. \quad \blacksquare$$

Exercise 3.2. *Prove that*

$$\left| \frac{1}{n} \sum_{i=1}^n \lfloor \frac{n}{i} \rfloor - \sum_{i=1}^n \frac{1}{i} \right| < 1.$$

3.2 Sperner's Theorem

Definition 3.3. *Let $\mathcal{F} \subseteq 2^{[n]}$ be a family of subsets of $[n]$. We say \mathcal{F} is **independent** (or \mathcal{F} is an **independent system**), if for any two $A, B \in \mathcal{F}$, we have $A \not\subseteq B$ and $B \not\subseteq A$. In other words, \mathcal{F} is independent if and only if there is no "containment" relationship between any two subsets of \mathcal{F} .*

Fact 3.4. *For a fixed $k \in [n]$, $\binom{[n]}{k}$ is an independent system.*

Theorem 3.5 (Sperner's Theorem). *For any independent system \mathcal{F} of $[n]$, we have*

$$|\mathcal{F}| \leq \binom{n}{\lfloor \frac{n}{2} \rfloor}.$$

First we define a chain.

Definition 3.6. A chain of subsets of $[n]$ is a sequence of distinct subsets such that

$$A_1 \subseteq A_2 \subseteq A_3 \subseteq \cdots \subseteq A_k.$$

First proof of Sperner's Theorem (Double-Counting). A maximal chain is a chain with the property that no other subsets of $[n]$ can be inserted into it to find a longer chain. We have the following observations.

(1). Any maximal chain looks like:

$$\phi \subseteq \{x_1\} \subseteq \{x_1, x_2\} \subseteq \cdots \subseteq \{x_1, \dots, x_k\} \subseteq \cdots \subseteq \{x_1, \dots, x_n\}.$$

(2). There are exactly $n!$ maximal chains.

This is because any such a maximal chain, say $\mathcal{C} : \phi \subseteq \{x_1\} \subseteq \{x_1, x_2\} \subseteq \cdots \subseteq \{x_1, x_2, \dots, x_n\}$, defines a unique permutation:

$$\pi : [n] \rightarrow [n], \pi(i) = x_i, \forall i \in [n].$$

Now we count the number of pairs (\mathcal{C}, A) satisfying that:

- \mathcal{C} is a maximal chain of $[n]$.
- $A \in \mathcal{C} \cap \mathcal{F}$.

Recall the rule of double counting given at the beginning that

$$\sum_{\mathcal{C}} N_{\mathcal{C}} = \text{the number of pairs } (\mathcal{C}, A) = \sum_A N_A,$$

where $N_{\mathcal{C}}$ is the number of subsets $A \in \mathcal{C} \cap \mathcal{F}$ and N_A is the number of maximal chains \mathcal{C} containing A . It is key to observe that

- $N_{\mathcal{C}} \leq 1$,
- $N_A = |A|!(n - |A|)!$

So we have

$$\begin{aligned} n! &= \sum_{\mathcal{C}} 1 \geq \sum_{\mathcal{C}} N_{\mathcal{C}} = \sum_{A \in \mathcal{F}} N_A = \sum_{A \in \mathcal{F}} |A|!(n - |A|)! \\ &= \sum_{A \in \mathcal{F}} \frac{n!}{\binom{n}{|A|}} \geq \sum_{A \in \mathcal{F}} \frac{n!}{\binom{n}{\lfloor \frac{n}{2} \rfloor}} = \frac{n!}{\binom{n}{\lfloor \frac{n}{2} \rfloor}} |\mathcal{F}|, \end{aligned}$$

which implies that

$$|\mathcal{F}| \leq \binom{n}{\lfloor \frac{n}{2} \rfloor}.$$

This finishes the proof. ■

Now we give another proof of Sperner's Theorem.

Definition 3.7. A chain is symmetric if it consists of subsets of sizes $k, k+1, \dots, \lfloor \frac{n}{2} \rfloor, \dots, n-k-1, n-k$ for some $k \geq 0$.

For example, when $n = 3$, $\{\{2\}, \{2, 3\}, \{1, 2, 3\}\}$ is not symmetric. And when $n = 4$, $\{\emptyset, \{1, 2, 3\}\}$ is not symmetric.

Theorem 3.8. The family $2^{[n]}$ can be partitioned into a disjoint union of symmetric chains.

First proof of Theorem 3.8. We prove by induction on n .

The base case is $n = 1$. The family $2^{[1]} = 2^{[1]} = \{\emptyset, \{1\}\}$, which itself is a symmetric chain. Thus this theorem is true for $n = 1$.

Now we may assume that $2^{[n]}$ can be partitioned into a disjoint union of symmetric chains e_1, e_2, \dots, e_t . Consider $2^{[n+1]}$, For any

$$e_i = \{P_k \subseteq P_{k+1} \subseteq \dots \subseteq P_{n-k}\},$$

define two new symmetric chains for $2^{[n+1]}$:

$$e'_i = \{P_{k+1} \subseteq P_{k+2} \subseteq \dots \subseteq P_{n-k}\},$$

and

$$e''_i = \{P_k \subseteq (P_k \cup \{n+1\}) \subseteq (P_{k+1} \cup \{n+1\}) \subseteq \dots \subseteq (P_{n-k} \cup \{n+1\})\}.$$

We assert that $\cup_i \{e'_i, e''_i\}$ is a disjoint union of symmetric chain for $2^{[n+1]}$. ■

Exercise 3.9. Prove that $\cup_i \{e'_i, e''_i\}$ is a disjoint union of symmetric chain for $2^{[n+1]}$.

Second proof of Theorem 3.8. For each $A \in 2^{[n]}$, we define a sequence “ $a_1 a_2 \dots a_n$ ” consisting of left and right parentheses by defining

$$a_i = \begin{cases} “(”, & \text{if } i \in A \\ “)”, & \text{otherwise.} \end{cases}$$

We then define the “partial pairing of parentheses” as follows:

- (1). First, we pair up all pairs “()” of adjoint parentheses.
- (2). Then, we delete these already paired parentheses.
- (3). Repeat the above process until nothing can be done.

Note that when this process stops, the remaining unpaired parentheses must look like this:

)))((((

We say two subsets $A, B \in 2^{[n]}$ have the same partial pairing, if the paired parentheses are the same (even in the same positions).

We can define an equivalence “ \sim ” on $2^{[n]}$ by letting $A \sim B$ if and only if A, B have the same partial pairing.

Exercise 3.10. Each equivalence class indeed forms a symmetric chain.

Using this fact, now we see that $2^{[n]}$ can be partitioned into disjoint equivalence classes, which are disjoint symmetric chains. This finishes the proof. ■

Theorem 3.8 can rapidly imply Sperner's Theorem.

Second proof of Sperner's Theorem. Note that by definition, any symmetric chain contains exactly one subset of size $\lfloor \frac{n}{2} \rfloor$. Since there are $\binom{n}{\lfloor \frac{n}{2} \rfloor}$ many subsets of size $\lfloor \frac{n}{2} \rfloor$, by Theorem 3.8, we see that any partition of $2^{[n]}$ into symmetric chains has to consist of exactly $\binom{n}{\lfloor \frac{n}{2} \rfloor}$ symmetric chains. Each symmetric chain can contain at most one subset from \mathcal{F} and thus we see $|\mathcal{F}| \leq \binom{n}{\lfloor \frac{n}{2} \rfloor}$. ■

3.3 Littlewood-Offord Problem

Theorem 3.11. Fix a vector $\vec{a} = (a_1, a_2, \dots, a_n)$ with each $|a_i| \geq 1$. Let $S = \{\vec{\epsilon} = (\epsilon_1, \epsilon_2, \dots, \epsilon_n) : \epsilon_i \in \{1, -1\} \text{ and } \vec{\epsilon} \cdot \vec{a} \in (-1, 1)\}$, then $|S| \leq \binom{n}{\lfloor \frac{n}{2} \rfloor}$.

Remark: Note that this is tight for many vectors \vec{a} .

Proof. For any $\vec{\epsilon} \in S$, define $A_{\vec{\epsilon}} = \{i \in [n] : a_i \epsilon_i > 0\}$. Let $\mathcal{F} = \{A_{\vec{\epsilon}} : \vec{\epsilon} \in S\}$. Then we have

$$|S| = |\mathcal{F}|.$$

Now we claim that \mathcal{F} is an independent system. Suppose for a contradiction that there exist $A_{\vec{\epsilon}_1}, A_{\vec{\epsilon}_2} \in \mathcal{F}$ with $A_{\vec{\epsilon}_1} \subseteq A_{\vec{\epsilon}_2}$. That also says,

$$\begin{cases} \vec{\epsilon}_1 \cdot \vec{a} \in (-1, 1), \\ \vec{\epsilon}_2 \cdot \vec{a} \in (-1, 1), \end{cases}$$

which imply that

$$|\epsilon_1 \cdot \vec{a} - \epsilon_2 \cdot \vec{a}| < 2.$$

By definition, we have

$$\vec{\epsilon}_1 \cdot \vec{a} = \sum_{i \in A_{\vec{\epsilon}_1}} |a_i| - \sum_{i \notin A_{\vec{\epsilon}_1}} |a_i| = 2 \sum_{i \in A_{\vec{\epsilon}_1}} |a_i| - \sum_{i=1}^n |a_i|.$$

Since $A_{\vec{\epsilon}_1} \subseteq A_{\vec{\epsilon}_2}$, we also have that

$$\vec{\epsilon}_2 \cdot \vec{a} - \vec{\epsilon}_1 \cdot \vec{a} = 2 \left(\sum_{i \in A_{\vec{\epsilon}_2}} |a_i| - \sum_{j \in A_{\vec{\epsilon}_1}} |a_j| \right) \geq 2|a_k| \geq 2, \text{ for some } k \in A_{\vec{\epsilon}_2} \setminus A_{\vec{\epsilon}_1}.$$

This is a contradiction. By Sperner's Theorem, we have $|S| = |\mathcal{F}| \leq \binom{n}{\lfloor \frac{n}{2} \rfloor}$. This finishes the proof. ■

3.4 Turán Type Problems

Definition 3.12. A graph G is **bipartite** if its vertex set can be partitioned into two parts (say A and B) such that each edge joints one vertex in A and another in B .

This is equivalent to say that $V(G)$ can be partitioned into two independent subsets. And we say (A, B) is a bipartition of G . For example, all even cycles C_{2k} are bipartite, while all odd cycles C_{2k+1} are not.

Definition 3.13. Let $K_{a,b}$ be the **complete bipartite** graph with two parts of sizes a and b . This is a bipartite graph with edge set $\{\{i, j\} : i \in A, j \in B\}$ where $|A| = a$ and $|B| = b$.

Definition 3.14. Given a graph H , we say a graph G is **H -free** if G dose not contain a copy of H as its subgraph.

For example, $K_{a,b}$ is K_3 -free.

Definition 3.15. For fixed graph H , let the **Turán number of H** , denoted by $\text{ex}(n, H)$, be the maximum number of edges in an n -vertex H -free graph G .

Theorem 3.16. $\text{ex}(n, C_4) \leq \frac{n}{4}(1 + \sqrt{4n - 3})$.

Proof. Let G be a C_4 -free graph with n vertices. We need to show that $e(G) \leq \frac{n}{4}(1 + \sqrt{4n - 3})$. Consider $S = \{(\{u_1, u_2\}, w) : u_1 w u_2 \text{ is a path of length 2 in } G\}$. Since G is C_4 -free, for fixed $\{u_1, u_2\}$, there is at most one vertex w such that $(\{u_1, u_2\}, w) \in S$. So we have

$$|S| = \sum_{\{u_1, u_2\}} \text{the number of } (\{u_1, u_2\}, w) \in S \leq \sum_{\{u_1, u_2\}} 1 = \binom{n}{2}.$$

On the other hand, fixed a vertex w , the number of $\{u_1, u_2\}$ such that $(\{u_1, u_2\}, w) \in S$ exactly equals $\binom{d(w)}{2}$, which implies that

$$|S| = \sum_{w \in V(G)} \binom{d(w)}{2} = \frac{1}{2} \sum_{w \in V(G)} d^2(w) - e(G).$$

Putting the above together, we have

$$\binom{n}{2} \geq |S| = \frac{1}{2} \sum_{w \in V(G)} d^2(w) - e(G).$$

Using Cauchy-Schwarz inequality, we have

$$\frac{n^2 - n}{2} \geq \frac{n}{2} \sum_{w \in V(G)} \frac{d^2(w)}{n} - e(G) \geq \frac{n}{2} \sum_{w \in V(G)} \left(\frac{d(w)}{n}\right)^2 - e(G),$$

which implies that

$$\frac{2e^2(G)}{n} - e(G) \leq \frac{n^2 - n}{2}.$$

Solving it, we can derive easily that $e(G) \leq \frac{n}{4}(1 + \sqrt{4n - 3})$. ■

Exercise 3.17. Prove that for all positive integer $n \geq 4$, $\text{ex}(n, C_4) < \frac{n}{4}(1 + \sqrt{4n-3})$.

Hint: Look up the Friendship Theorem.

Corollary 3.18. We have $\text{ex}(n, C_4) \leq (\frac{1}{2} + o(n))n^{\frac{3}{2}}$, where $o(n) \rightarrow 0$ as $n \rightarrow \infty$.

The upper bound in Corollary 3.18 is asymptotically tight because there is a construction as follows.

Let p be a prime. Let

$$V = (\mathbb{Z}_p \setminus \{0\}) \times \mathbb{Z}_p$$

and

$$E = \{(a, b), (c, d)\} : a, c \in \mathbb{Z}_p \setminus \{0\}, b, d \in \mathbb{Z}_p \text{ and } ac = b + d\}.$$

We have $|V| = (p-1)p$ and $d((a, b)) = p-1$, for any $(a, b) \in V$. Thus we have $|E| = \frac{(p-1)^2 p}{2} \sim \frac{|V|^{\frac{3}{2}}}{2}$. Finally we explain that $G = (V, E)$ is C_4 -free. For any $(a_1, b_1), (a_2, b_2) \in V$, if there exist a vertex (say (c, d)) which is their common neighbour, (c, d) satisfies the following condition:

$$\begin{cases} a_1 c = b_1 + d \\ a_2 c = b_2 + d. \end{cases}$$

There is no multiple solution of this equation.

Theorem 3.19 (Kővári-Sós-Turán Theorem).

$$\text{ex}(n, K_{s,t}) \leq \frac{1}{2}(t-1)^{\frac{1}{s}} n^{2-\frac{1}{s}} + \frac{1}{2}(s-1)n$$

for all $t, s \geq 2$.

Proof. Let G be an n -vertices $K_{s,t}$ free graph with $e(G) \geq \frac{1}{2}sn$ (otherwise we are done). We aim to show $e(G) \leq \frac{1}{2}(t-1)^{\frac{1}{s}} n^{2-\frac{1}{s}} + \frac{1}{2}(s-1)n$. We count the number T of s -stars $K_{1,s}$ as follows. On one hand, $T = \sum_{w \in V(G)} \binom{d(w)}{s}$. On the other hand, $T \leq (t-1) \binom{n}{s}$.

We define

$$f(x) = \begin{cases} 0 & , \text{ if } x < s, \\ \binom{x}{s} & , \text{ if } x \geq s. \end{cases}$$

When $x \geq 0$, $f(x)$ is a convex function. Let $d = \frac{2e(G)}{n}$, by Jensen's inequality,

$$\frac{(t-1) \binom{n}{s}}{n} \geq \frac{T}{n} = \frac{1}{n} \sum_w f(d(w)) \geq f\left(\frac{\sum_w d(w)}{n}\right) = f\left(\frac{2e(G)}{n}\right) \geq \frac{(d-s+1)^s}{s!}.$$

Thus

$$d \leq ((t-1)(n-1)(n-2) \dots (n-s+1))^{\frac{1}{s}} + (s-1) \leq (t-1)^{\frac{1}{s}} n^{1-\frac{1}{s}} + (s-1).$$

Then we have

$$e(G) = \frac{nd}{2} \leq \frac{1}{2}(t-1)^{\frac{1}{s}} n^{2-\frac{1}{s}} + \frac{1}{2}(s-1)n,$$

finishing the proof. ■

3.5 Sperner's Lemma

Let us consider the following application of Corollary 2.8. First we draw a triangle in the plane, with 3 vertices $A_1A_2A_3$. Then we divide this triangle $\Delta = A_1A_2A_3$ into small triangles such that no triangle can have a vertex inside an edge of any other small triangle. Then we assign 3 colors (say 1,2,3) to all vertices of these triangles, under the following rules.

- (1) The vertex A_i is assigned by color i for $i \in [3]$.
- (2) All vertices lying on the edge A_iA_j of the large triangle are assigned by the color i or j .
- (3) All interior vertices are assigned by any color 1,2,3.

Lemma 3.20 (Sperner's Lemma (a planar version)). *For any assignment of colors described as above, there always exists a small triangle whose three vertices are assigned by three colors 1, 2, 3.*

Proof. Define an auxiliary graph G as follows.

- Its vertices are the faces of small triangles and the outer face. Let z be the vertex representing the outer face.
- Two vertices of G are adjacent, if the two corresponding faces are neighboring faces and the two endpoints of their common edge are colored by 1 and 2.

We consider the degree of any vertex $v \in V(G) \setminus \{z\}$.

- (1) If the face of v has NO two endpoints with color 1 and 2, then $d_G(v) = 0$.
- (2) If the face of v has 2 endpoints with color 1 and 2, then let k be the color of the third endpoint of this face. If $k \in \{1, 2\}$, then $d_G(v) = 2$. Otherwise $k = 3$, then $d_G(v) = 1$ and the vertices of this triangle are assigned by three different colors 1,2,3.

Thus we have that $d_G(v)$ is odd if and only if $d_G(v) = 1$, and then the face of v has colors 1,2,3. Now we consider $d_G(z)$ and claim that it must be odd. Indeed, the edge of G incident to z obviously have to go across A_1A_2 . Consider the sequence of the colors of the endpoints on A_1A_2 , from A_1 to A_2 . Then $d_G(z)$ equals the number of alternations between 1 and 2 in this sequence. It is easy to check that $d_G(z)$ must be odd. By Corollary 2.8, since the graph G has a vertex z with odd degree, there must be another vertex $v \in V(G) \setminus \{z\}$ with odd degree. Then $d(v) = 1$ and the face of v has colors 1,2,3. ■


Before we introduce an interesting application of Sperner's lemma, we introduce the following theorem first.

Theorem 3.21 (One-dimensional fixed point theorem). *For any continuous function $f : [0, 1] \rightarrow [0, 1]$, there exists a point $x \in [0, 1]$ such that $f(x) = x$.*

Such an x is called a fixed point of the function f . The theorem can be proved by considering the function $g(x) = f(x) - x$. This is a continuous function with $g(0) \geq 0$ and $g(1) \leq 0$. Intuitively it is quite clear that the graph of such a continuous function can not jump across the x -axis and therefore it has to intersect it, and hence g is 0 at some point of $[0, 1]$. Prove the existence of such a point rigorously requires quite some work. In analysis, this result appears under heading "Darboux theorem".

If we replace the 1-dimensional interval from the Theorem 3.21 by a triangle in the plane, or by a tetrahedron in the 3-dimensional space, or by their analogs in higher dimensions, we will have Brouwer's fixed point theorem. Here we prove only the 2-dimensional version by Spener's lemma.

Let Δ denote a triangle in the plane. For simplicity, let us take the triangle with vertices $A_1 = (1, 0)$, $A_2 = (0, 1)$, and $A_3 = (0, 0)$:



Theorem 3.22 (Brouwer's Fixed Point theorem in 2-dimension). *Every continuous function $f : \Delta \rightarrow \Delta$ has a fixed point x , that is, $f(x) = x$.*

Proof. Define three auxiliary functions $\beta_i : \Delta \rightarrow R$ for $i \in \{1, 2, 3\}$ as follows:


For any $a = (x, y) \in \Delta$,

$$\begin{cases} \beta_1(a) = x, \\ \beta_2(a) = y, \\ \beta_3(a) = 1 - x - y. \end{cases}$$

For any continuous $f : \Delta \rightarrow \Delta$, define $M_i = \{a \in \Delta : \beta_i(a) \geq \beta_i(f(a))\}$ for $i \in \{1, 2, 3\}$. Then we have the following facts.

- (1) Any point $a \in \Delta$ belongs to at least one M_i .
- (2) If $a \in M_1 \cap M_2 \cap M_3$, then a is a fixed point.

Consider a sequence of refinements $\{\Delta_1, \Delta_2, \dots\}$ of Δ such that the maximum diameter of small triangles in Δ_n is going to 0 as $n \rightarrow +\infty$. For example, we can consider the refining triangulations of the triangle Δ as follows:



We want to define a coloring $\phi : \Delta \rightarrow \{1, 2, 3\}$ such that

- (a) Any $a \in \Delta$ with $\phi(a) = i$ belongs to M_i .
- (b) The coloring ϕ satisfies the conditions of Spener's Lemma for any subdivision Δ_n of Δ .

Next we show such ϕ exists. This is because

- For the point A_i (say $i = 1$), we have that $A_1 = (1, 0) \in M_1$, so we can let $\phi(A_i) = i$;
- Consider a vertex $a = (x, y) \in A_1A_2$, i.e., $x + y = 1$. Since $\beta_1(f(a)) + \beta_2(f(a)) \leq 1 = x + y = \beta_1(a) + \beta_2(a)$, so we must have at least one of $\beta_1(f(a)) \leq \beta_1(a)$ and $\beta_2(f(a)) \leq \beta_2(a)$ holds, which means that $a \in M_1 \cup M_2$.

Applying Sperner's Lemma to each Δ_n and the coloring ϕ , we get that there exists a small triangle $A_1^{(n)}A_2^{(n)}A_3^{(n)}$ in Δ_n which has three different colors 1,2,3.

Consider the sequence $\{A_1^{(n)}\}_{n \geq 1}$. Since everything is bounded, there is a subsequence $\{A_1^{(n_k)}\}_{k \geq 1}$ such that $\lim_{k \rightarrow +\infty} A_1^{(n_k)} = p \in \Delta$ exists. Since the diameter of $A_1^{(n)}A_2^{(n)}A_3^{(n)}$ is going to be 0 as $n \rightarrow +\infty$, we see that $\lim_{k \rightarrow +\infty} A_2^{(n_k)} = \lim_{k \rightarrow +\infty} A_3^{(n_k)} = p$. Since $\beta_i(A_i^{(n_k)}) \geq \beta_i(f(A_i^{(n_k)}))$ for $i \in [3]$ and f is continuous, we get $\beta_i(p) = \lim_{k \rightarrow +\infty} \beta_i(A_i^{(n_k)}) \geq \lim_{k \rightarrow +\infty} \beta_i(f(A_i^{(n_k)})) = \beta_i(f(p))$ for $i \in [3]$. This implies that $p \in M_1 \cap M_2 \cap M_3$, so p is a fixed point of f , that is, $f(p) = p$. ■